

STRENGTH OF CONVERGENCE IN THE ORBIT SPACE OF A TRANSFORMATION GROUP

ROBERT ARCHBOLD AND ASTRID AN HUEF

ABSTRACT. Let (G, X) be a second-countable transformation group with G acting freely on X . It is shown that measure-theoretic accumulation of the action and topological strength of convergence in the orbit space X/G provide equivalent ways of quantifying the extent of non-properness of the action. These notions are linked via the representation theory of the transformation-group C^* -algebra $C_0(X) \rtimes G$.

1. INTRODUCTION

Let G be a locally compact Hausdorff group acting continuously on a locally compact Hausdorff space X , so that (G, X) is a transformation group. In [11] Green gave a seminal example of a non-proper action of $G = \mathbb{R}$ on a subspace X of $\mathbb{R}^3$; this example was generalised in [18] by Rieffel who replaced the double folding of orbits by variable folding using a repetition number associated to each orbit. The main purpose of this paper is to show that, for free actions, the fundamental measure-theoretic and topological properties of these examples persist into the general case. In particular, we show that the local accumulation of Haar measure (which arises from the folding of orbits in the case of the Green-Rieffel examples) corresponds exactly to a topological notion of strength of convergence in the orbit space X/G from [2]. The latter notion is motivated by the phenomenon of strength of convergence for the Kirillov orbits associated to a nilpotent Lie group [15, 4].

To show that these notions of “counting by measure” and “counting by topology” give equivalent ways of quantifying the extent of non-properness of an action, we use the transformation-group C^* -algebra $C_0(X) \rtimes G$. It is well-known that the representation theory of $C_0(X) \rtimes G$ is closely related to the properties of the action of G on X , particularly when G and X are second countable. In [19, Theorem 3.1] Williams showed that $C_0(X) \rtimes G$ is liminal (that is, the image of every irreducible representation π of $C_0(X) \rtimes G$ is the compact operators) if and only if each orbit is closed in X and all the stability subgroups are liminal. More generally, Gootman proved in [10, Theorem 3.3] that $C_0(X) \rtimes G$ is Type I if and only if each orbit is locally closed in X and each stability subgroup is Type I. Further, if $C_0(X) \rtimes G$ is Type I and the action of G on X is free, the spectrum $(C_0(X) \rtimes G)^\wedge$ of $C_0(X) \rtimes G$ is homeomorphic to X/G [11, Lemma 16].

Date: August 18, 2005.

2000 Mathematics Subject Classification. 22D30, 46L05, 46L30, 46L55, 54H15, 57S05.

Key words and phrases. Transformation group, orbit space, k -times convergence, spectrum of a C^* -algebra, multiplicity of a representation, trace function.

This research was supported by grants from the Royal Society and the University of New South Wales.

A subset K of X is *wandering* if the set

$$\{s \in G : s \cdot K \cap K \neq \emptyset\}$$

is relatively compact in G . The action of G on X is called *proper* if every compact subset of X is wandering, or equivalently, if the map

$$(s, x) \mapsto (s \cdot x, x) : G \times X \rightarrow X \times X$$

is proper in the sense that inverse images of compact subsets are compact. If every point of X has a wandering neighbourhood, then X is a *Cartan G -space* [16, Definition 1.1.2]. For free actions, the conditions on the spectrum of $C_0(X) \rtimes G$ corresponding to proper actions and Cartan G -spaces are that $C_0(X) \rtimes G$ has continuous trace [11, Theorem 17] and $C_0(X) \rtimes G$ is a Fell algebra [12], respectively. That the action of G on X is proper if and only if the orbit space X/G is Hausdorff and X is a Cartan G -space [16, Theorem 1.2.9] is thus closely related to a C^* -algebra having continuous trace if and only if it is a Fell algebra with Hausdorff spectrum [8, §4.5].

More general than a Cartan G -space is the notion of an integrable action of G on $C_0(X)$ from [18, Definition 1.10]; after translating that definition to the transformation group (G, X) , an action of G on X is *integrable* if and only if, for each compact subset N of X ,

$$\sup_{x \in N} \nu(\{s \in G : s \cdot x \in N\}) < \infty, \quad (1.1)$$

where ν is a (right) Haar measure on G . For free actions, it is shown in [13, Theorem 4.8] that the action is integrable if and only if $C_0(X) \rtimes G$ has bounded trace (that is, there is a dense ideal J of $C_0(X) \rtimes G$ such that $\pi \mapsto \text{tr}(\pi(a))$ is bounded on $(C_0(X) \rtimes G)^\wedge$ for every fixed positive element a of J). On the other hand, it follows from [5, Theorem 2.6] that $C_0(X) \rtimes G$ has bounded trace if and only if the upper multiplicity $M_U(\pi)$ is finite for every irreducible representation π of $C_0(X) \rtimes G$ (see §2 for the definition of the multiplicity numbers). So, when the action is free and integrable, it is natural to ask how the finite upper multiplicity numbers of the representations of $C_0(X) \rtimes G$ are related to the finite suprema occurring in (1.1).

Suppose that $(x_n)_{n \geq 1}$ is a sequence in X converging to $z \in X$, and that $\text{Ind } \epsilon_{x_n}$ and $\text{Ind } \epsilon_z$ are the corresponding induced representations of $C_0(X) \rtimes G$. The examples in [11, 18, 7, 13] suggest that the upper and lower multiplicities $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n}))$ and $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n}))$ of $\text{Ind } \epsilon_z$ relative to the sequence $(\text{Ind } \epsilon_{x_n})_{n \geq 1}$ depend on the ratios

$$\frac{\nu(\{s : s \cdot x_n \in V\})}{\nu(\{s : s \cdot z \in V\})} \quad (1.2)$$

which compare the amount of “time” x_n and z spend in a neighbourhood V of z under the action of G .

By contrast, a topological rather than measure-theoretic approach was adopted in [2]. Motivated by the notion of strength of convergence for the Kirillov orbits associated to a nilpotent Lie group (see [15, Definition 2.10] and [4, Theorem 2.4]), a sequence $(x_n)_{n \geq 1}$ in X is said to *converge k -times in X/G to $z \in X$* if there exist k sequences $(t_n^{(1)})_n, (t_n^{(2)})_n, \dots, (t_n^{(k)})_n$ in G such that

- (1) $t_n^{(i)} \cdot x_n \rightarrow z$ as $n \rightarrow \infty$ for $1 \leq i \leq k$, and
- (2) if $1 \leq i < j \leq k$ then $t_n^{(j)}(t_n^{(i)})^{-1} \rightarrow \infty$ as $n \rightarrow \infty$.

It follows from [2, Theorem 2.3] that if $(x_n)_{n \geq 1}$ converges k -times in X/G to z then $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \geq k$.

The main theorem of this paper shows that, for free actions, the measure-theoretic and topological approaches are equivalent. Of the five equivalent conditions of the theorem, only the second involves the transformation-group C^* -algebra $C_0(X) \rtimes G$; its representation theory, via the multiplicity numbers, is the key to passing from topological convergence to measure accumulation.

Theorem 1.1. *Suppose that (G, X) is a free second-countable transformation group. Let k be a positive integer, let $z \in X$ and let $(x_n)_{n \geq 1}$ be a sequence in X . Assume that $G \cdot z$ is locally closed in X . Then the following are equivalent:*

- (1) *the sequence $(x_n)_n$ converges k -times in X/G to z ;*
- (2) $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \geq k$;
- (3) *for every open neighbourhood V of z such that $\{s \in G : s \cdot z \in V\}$ is relatively compact we have*

$$\liminf_n \nu(\{s \in G : s \cdot x_n \in V\}) \geq k \nu(\{s \in G : s \cdot z \in V\});$$

- (4) *there exists a real number $R > k - 1$ such that for every open neighbourhood V of z with $\{s \in G : s \cdot z \in V\}$ relatively compact we have*

$$\liminf_n \nu(\{s \in G : s \cdot x_n \in V\}) \geq R \nu(\{s \in G : s \cdot z \in V\});$$

- (5) *there exists a decreasing sequence of basic compact neighbourhoods $(W_m)_{m \geq 1}$ of z such that, for each $m \geq 1$,*

$$\liminf_n \nu(\{s \in G : s \cdot x_n \in W_m\}) > (k - 1) \nu(\{s \in G : s \cdot z \in W_m\}).$$

In (3)–(5) above, it is to be understood that the limit infima are calculated in $[0, \infty]$ and that some of the values $\nu(\{s \in G : s \cdot x_n \in V\})$ might be infinite. The equivalence of (3), (4) and (5) demonstrates a surprising emergence of integer values from what appears in general to be a continuous setting. Note that if the action is proper, or more generally if X is a Cartan G -space, then the upper multiplicity $M_U(\text{Ind } \epsilon_z) = 1$ for all $z \in X$ [11, 12, 1], and hence there can only be 1-times convergence in X/G to z .

This paper is organised as follows. In §2 we establish our conventions and discuss our hypotheses. In §3 we establish a relationship between the ratios (1.2) and the lower and upper multiplicities $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n}))$ and $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n}))$ relative to the sequence $(\text{Ind } \epsilon_{x_n})_n$ (see Theorems 3.5 and 3.6). In §4 we link the limit suprema and infima of measure ratios and topological strength of convergence. A crucial ingredient for the results in §3 and §4 is a new technical lemma (Lemma 3.2) which shows that a key property which we have observed in Green's example [11] can be abstracted to the general situation.

We combine our results in §5 to prove Theorem 1.1. The first corollary of Theorem 1.1 is the analogous result involving k -times convergence of a subsequence of $(x_n)_n$, the upper multiplicity $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n}))$ and limit suprema of the measure ratios (Theorem 5.3).

We apply our results to examples in §6. In particular, we consider Rieffel's example of a free transformation group [18, Example 1.18] where the orbit space X/G consists of a sequence $(G \cdot x_n)_{n \geq 1}$ converging to $G \cdot x_0$. For each $n \geq 1$ the orbit has $L_n + 1$ folds,

where L_n is a repetition number. We show that $M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = \liminf (L_n + 1)$ and $M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = \limsup (L_n + 1)$.

In Appendix A we establish sequence versions of [6, Propositions 2.2 and 2.3] which are needed in §3 and in §5.

2. PRELIMINARIES

Throughout, (G, X) is a locally compact Hausdorff transformation group: thus G is a locally compact Hausdorff group and X is a locally compact Hausdorff space together with a jointly continuous map $(s, x) \mapsto s \cdot x$ from $G \times X$ to X such that $s \cdot (t \cdot x) = st \cdot x$ and $e \cdot x = x$. In all our main results the action is assumed to be *free*, that is $s \cdot x = x$ implies $s = e$. We also assume that both G and X are second countable, and hence both are normal spaces [14, Chapter 4, Lemma 1].

The action of G on X lifts to a strongly continuous action lt of G by automorphisms of $C_0(X)$ given by $\text{lt}_s(f)(x) = f(s^{-1} \cdot x)$. The associated transformation-group C^* -algebra $C_0(X) \rtimes G$ is the C^* -algebra which is universal for the covariant representations of the C^* -dynamical system $(C_0(X), G, \text{lt})$. More concretely, $C_0(X) \rtimes G$ is the enveloping C^* -algebra of the Banach $*$ -algebra $L^1(G, C_0(X))$ of functions $f : G \rightarrow C_0(X)$ which are integrable with respect to a fixed left Haar measure μ on G , with multiplication and involution given by

$$f * g(s) := \int_G f(r) \text{lt}_r(g(r^{-1}s)) d\mu(r) \quad \text{and} \quad f^*(s) := \Delta(s^{-1}) \text{lt}_s(f(s^{-1}))^*,$$

where Δ is the modular function associated with μ .

Suppose that G acts freely on X and let ν be the right Haar measure such that $\nu(E) = \mu(E^{-1})$. As in [19, 20, 13], for each $x \in X$ we realise the induced representation $\text{Ind } \epsilon_x$ (where $\epsilon_x : C_0(X) \rightarrow \mathbb{C}$ is evaluation at x) on the Hilbert space $L^2(G, \nu)$ as the integrated form

$$\text{Ind } \epsilon_x = \tilde{\epsilon}_x \times \lambda,$$

where $(\tilde{\epsilon}_x(f)\xi)(s) = f(s \cdot x)\xi(s)$ and $(\lambda_t\xi)(s) = \Delta(t)^{1/2}\xi(t^{-1}s)$ for $f \in C_0(X)$, $\xi \in L^2(G, \nu)$ and $r, s \in G$ (see [19, Lemma 4.14]). Thus

$$\begin{aligned} (\text{Ind } \epsilon_x(f)\xi)(s) &= \int_G f(t, s \cdot x)\xi(t^{-1}s)\Delta(t)^{1/2} d\mu(t) \\ &= \int_G f(sw^{-1}, s \cdot x)\Delta(sw^{-1})^{1/2}\xi(w) d\nu(w) \end{aligned} \quad (2.1)$$

for $f \in C_c(G, X)$ and $\xi \in L^2(G, \nu)$. Note that $\text{Ind } \epsilon_x$ is irreducible by, for example, [19, Proposition 4.2].

Let A be a C^* -algebra and let π be an irreducible representation of A ; we use the same symbol π to denote the unitary equivalence class of π in the spectrum $\hat{A}$ of A . If π_1 and π_2 are equivalent irreducible representations then $\text{tr}(\pi_1(a)) = \text{tr}(\pi_2(a))$ for all a in the positive cone A^+ of A , and so we may write unambiguously the expression $\text{tr}(\pi(a))$ whenever $\pi \in \hat{A}$ and $a \in A^+$.

We now recall the definitions of upper and lower multiplicity from [1], [6]. Unless stated otherwise, we shall always regard the Banach dual A^* as being equipped with the weak*-topology. We denote by $\mathcal{N}$ the weak*-neighbourhood base at zero in A^* consisting of all

open sets of the form

$$N = \{\psi \in A^* : |\psi(a_i)| < \epsilon, 1 \leq i \leq n\}$$

where $\epsilon > 0$ and $a_1, \dots, a_n \in A$. Let $P(A)$ be the set of pure states of A and let $\theta : P(A) \rightarrow \hat{A}$ be the continuous, open mapping given by $\theta(\phi) = \pi_\phi$ where π_ϕ is the GNS representation associated with ϕ [8, 3.4.11].

We write $\mathbb{P} = \mathbb{N} \setminus \{0\}$. Let $\pi \in \hat{A}$. First of all, we give descriptions of the upper and lower multiplicities $M_U(\pi)$ and $M_L(\pi)$. Let ϕ be a pure state of A associated with π and let $N \in \mathcal{N}$. Let

$$V(\phi, N) = \theta((\phi + N) \cap P(A)),$$

an open neighbourhood of π in $\hat{A}$. For $\sigma \in \hat{A}$ let $\mathcal{H}_\sigma$ be the Hilbert space of σ , and let

$$\text{Vec}(\sigma, \phi, N) = \{\eta \in \mathcal{H}_\sigma : \|\eta\| = 1, \langle \sigma(\cdot)\eta, \eta \rangle \in \phi + N\}.$$

Note that $\text{Vec}(\sigma, \phi, N)$ is nonempty if and only if $\sigma \in V(\phi, N)$. For $\sigma \in V(\phi, N)$ we define $d(\sigma, \phi, N)$ to be the supremum (in $\mathbb{P} \cup \{\infty\}$) of the cardinalities of finite orthonormal subsets of $\text{Vec}(\sigma, \phi, N)$. It is convenient to define $d(\sigma, \phi, N) = 0$ for $\sigma \in \hat{A} \setminus V(\phi, N)$.

From [1, §2 and Proposition 3.4] we have

$$M_U(\pi) = \inf_{N \in \mathcal{N}} \left(\limsup_{\sigma \rightarrow \pi} d(\sigma, \phi, N) \right) \in \mathbb{P} \cup \{\infty\}$$

and, if π is not open in $\hat{A}$,

$$M_L(\pi) = \inf_{N \in \mathcal{N}} \left(\liminf_{\sigma \rightarrow \pi, \sigma \neq \pi} d(\sigma, \phi, N) \right) \in \mathbb{P} \cup \{\infty\}.$$

As noted in [1, Lemma 2.1], $M_U(\pi)$ and $M_L(\pi)$ are independent of the choice of ϕ . Elementary examples which motivate these definitions and illustrate the computations are given in [1, §2].

Now suppose, in addition, that $\Omega = (\pi_\alpha)_{\alpha \in \Lambda}$ is a net in $\hat{A}$. For $N \in \mathcal{N}$ let

$$M_U(\phi, N, \Omega) = \limsup_{\alpha} d(\pi_\alpha, \phi, N) \in \mathbb{N} \cup \{\infty\}.$$

Note that if $N' \in \mathcal{N}$ and $N' \subset N$ then $M_U(\phi, N', \Omega) \leq M_U(\phi, N, \Omega)$. We define

$$M_U(\pi, \Omega) = \inf_{N \in \mathcal{N}} M_U(\phi, N, \Omega) \in \mathbb{N} \cup \{\infty\}$$

(which is independent of the choice of ϕ by an argument similar to that used in the proof of [1, Lemma 2.1]). Similarly, for $N \in \mathcal{N}$, let

$$M_L(\phi, N, \Omega) = \liminf_{\alpha} d(\pi_\alpha, \phi, N) \in \mathbb{N} \cup \{\infty\}.$$

Then $M_L(\phi, N, \Omega)$ decreases with N and we define

$$M_L(\pi, \Omega) = \inf_{N \in \mathcal{N}} M_L(\phi, N, \Omega) \in \mathbb{N} \cup \{\infty\}$$

(which is, again, independent of the choice of ϕ). Note that it is not required that Ω converge to π . However it follows from these definitions that $M_U(\pi, \Omega) > 0$ if and only if π is a cluster point of Ω , and that $M_L(\pi, \Omega) > 0$ if and only if Ω converges to π .

We have the inequalities

$$M_L(\pi, \Omega) \leq M_U(\pi, \Omega) \leq M_U(\pi)$$

and, if Ω is convergent to π but eventually $\pi_\alpha \neq \pi$,

$$M_L(\pi) \leq M_L(\pi, \Omega)$$

(see [6, Proposition 2.1]). Also, if Ω_0 is a subnet of Ω then

$$M_L(\pi, \Omega) \leq M_L(\pi, \Omega_0) \leq M_U(\pi, \Omega_0) \leq M_U(\pi, \Omega).$$

Remark on hypotheses. In our major results, we assume that the action of G on X is free. Typically we focus on a fixed $z \in X$ together with a sequence $(x_n)_{n \geq 1}$ in X such that the orbit $G \cdot z := \{s \cdot z : s \in G\}$ is locally closed in X (in the sense that $G \cdot z$ is relatively open in its closure) and $G \cdot x_n \rightarrow G \cdot z$ in the orbit space X/G .

Lemma 2.1 explains the hypotheses in our main Theorem 1.1. In particular, if $G \cdot z$ is not locally closed then one cannot form the measure ratios (1.2). For each $x \in X$, let $\phi_x : G \rightarrow G \cdot x$ be the map $s \mapsto s \cdot x$.

Lemma 2.1. *Let (G, X) be a second-countable transformation group and ν a right Haar measure on G . Let $z \in X$ and suppose that the stability subgroup $S_z := \{s \in G : s \cdot z = z\}$ is compact. Then the following are equivalent:*

- (1) *the orbit $G \cdot z$ is not locally closed in X ;*
- (2) *for every $k \in \mathbb{P}$, the sequence $z, z, z, \dots$ converges k -times in X/G to z ;*
- (3) *for every open neighbourhood V of z , $\nu(\phi_z^{-1}(V)) = \infty$;*
- (4) *for every open neighbourhood V of z , $\phi_z^{-1}(V)$ is not relatively compact in G .*

Proof. Let $(V_n)_{n \geq 1}$ be a decreasing sequence of basic open neighbourhoods of z in X and let $(K_n)_{n \geq 1}$ be an increasing sequence of compact subsets of G such that $G = \bigcup_{n \geq 1} \text{Int}(K_n)$.

(1) $\implies$ (2). Suppose that $G \cdot z$ is not locally closed. Then $W \cap (\overline{G \cdot z} \setminus G \cdot z) \neq \emptyset$ for every neighbourhood W of z . Let $k \geq 1$. We will construct k sequences $(t_n^{(i)})_{n \geq 1} \subset G$, where $1 \leq i \leq k$, such that $t_n^{(i)} \cdot z \in V_n$ and $t_n^{(j)}(t_n^{(i)})^{-1} \notin K_n$ for each $n \geq 1$.

Temporarily fixing n , we construct $t_n^{(i)}$ as follows. Let $t_n^{(1)} = e$. Since $G \cdot z$ is not locally closed there exists $y \in V_n \cap (\overline{G \cdot z} \setminus G \cdot z)$. Since y is in the closure of $G \cdot z$ and V_n is open, given any compact subset K of G there exists $t_K \in G \setminus K$ such that $t_K \cdot z \in V_n$. So there exists $t_n^{(2)} \in G \setminus K_n$ such that $t_n^{(2)} \cdot z \in V_n$. Proceeding inductively we obtain $t_n^{(2)}, t_n^{(3)}, \dots, t_n^{(k)}$ such that

$$t_n^{(j)} \cdot z \in V_n \quad \text{and} \quad t_n^{(j)} \in G \setminus \left(\bigcup_{i=1}^{j-1} K_n t_n^{(i)} \right)$$

for $2 \leq j \leq k$. Letting n run, it follows easily from the properties of $(V_n)_{n \geq 1}$ and $(K_n)_{n \geq 1}$ that $z, z, z, \dots$ converges k -times in X/G to z .

(2) $\implies$ (3). Suppose that (2) holds. Let V be an open neighbourhood of z and $M > 0$. By the continuity of the action on the locally compact Hausdorff space X , there exists an open neighbourhood U of z and a compact neighbourhood K of e in G such that $K \cdot U \subset V$. Choose $k \in \mathbb{P}$ such that $k\nu(K) > M$. By (2) there exist k sequences $(t_n^{(i)})_{n \geq 1}$ such that $t_n^{(i)} \cdot z \rightarrow z$ as $n \rightarrow \infty$ for each $1 \leq i \leq k$, and

$$t_n^{(j)}(t_n^{(i)})^{-1} \rightarrow \infty \text{ as } n \rightarrow \infty \quad (1 \leq i < j \leq k).$$

Hence there exists n_0 such that $t_{n_0}^{(i)} \cdot z \in U$ for $1 \leq i \leq k$ and $t_{n_0}^{(j)}(t_{n_0}^{(i)})^{-1} \in G \setminus (K^{-1}K)$ for $1 \leq i < j \leq k$. Then $K t_{n_0}^{(i)} \cdot z \subset K \cdot U \subset V$ for $1 \leq i \leq k$ and $K t_{n_0}^{(i)} \cap K t_{n_0}^{(j)} = \emptyset$ unless $i = j$, and hence $\nu(\phi_z^{-1}(V)) \geq k\nu(K) > M$. Since M was arbitrary, the result follows.

(3) $\implies$ (4). Compact subsets have finite Haar measure, so this is immediate.

(4) $\implies$ (1). Suppose that $G \cdot z$ is locally closed in X . Then $G \cdot z$ is a relatively open subset of the locally compact space $\overline{G \cdot z}$ and hence $G \cdot z$ is locally compact. Thus $(G, G \cdot z)$ is a second-countable locally compact Hausdorff transformation group. In particular, it follows from [9, Theorem 1] that the map $\psi_z : G/S_z \rightarrow G \cdot z : sS_z \mapsto s \cdot z$ is a homeomorphism. Let U be an open subset of X such that $U \cap \overline{G \cdot z} = G \cdot z$. Let N be a compact neighbourhood of z in X such that $N \subset U$. Then $N \cap \overline{G \cdot z} = N \cap G \cdot z$ is a compact subset of $G \cdot z$. Hence $\psi_z^{-1}(N)$ is compact in G/S_z . Since S_z is compact, the quotient map $q_z : G \rightarrow G/S_z$ is proper, and hence $\phi_z^{-1}(V) = q_z^{-1}(\psi_z^{-1}(V))$ is compact in X . Let $V = \text{Int } N$; then $\phi_z^{-1}(V)$ is relatively compact in G . $\square$

If $M_U(\text{Ind } \epsilon_z)$ were finite, then we could assume, without loss of generality, that all orbits are closed in X . To see this, note that $\text{Ind } \epsilon_z$ restricts to an irreducible representation of the bounded-trace ideal J of $C_0(X) \times G$ [5, Theorem 2.8], and J is canonically isomorphic to $C_0(Y) \times G$ for some G -invariant open subset Y of X [13, Theorem 5.8]. Since J (and hence $C_0(Y) \times G$) is liminal, the orbits in Y are relatively closed in Y . By replacing X with Y we are then in the situation of closed orbits without changing any dynamical or representational properties at or near z ; in particular, the multiplicities for $\text{Ind } \epsilon_z$ are the same whether we compute them in the ideal $C_0(Y) \times G$ or in $C_0(X) \times G$ [5, Proposition 5.3]. However, we are also interested in examples where $M_U(\text{Ind } \epsilon_z) = \infty$ but the lower multiplicity $M_L(\text{Ind } \epsilon_z)$ is finite, and in examples where $M_L(\text{Ind } \epsilon_z) = M_U(\text{Ind } \epsilon_z) = \infty$.

3. MEASURE RATIOS AND BOUNDS ON MULTIPLICITY NUMBERS

We will frequently use the following well-known properties of a locally compact Hausdorff space X (see, for example, [14, Chapter 5, Theorem 18]): if U is a neighbourhood of a compact subset N of X , then U contains a compact neighbourhood V of N and there is a continuous function $f : X \rightarrow [0, 1]$ which is 1 on N and 0 on $X \setminus V$ (and hence on $X \setminus U$).

For $x \in X$ and $f \in C_0(X)$ we define the function $f_x : G \rightarrow \mathbb{C}$ by $f_x(s) = (f \circ \phi_x)(s) = f(s \cdot x)$.

Theorem 3.1. *Suppose that (G, X) is a free second-countable transformation group. Let $M \in \mathbb{R}$ with $M \geq 1$, let $z \in X$ and let $(x_n)_{n \geq 1}$ be a sequence in X . Assume that $G \cdot z$ is locally closed in X . Suppose that there exists an open neighbourhood V of z in X such that $\phi_z^{-1}(V)$ is relatively compact and*

$$\nu(\phi_{x_n}^{-1}(V)) \leq M\nu(\phi_z^{-1}(V))$$

frequently. Then $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \leq \lfloor M^2 \rfloor$.

Proof. Fix $\epsilon > 0$ such that $M^2(1+\epsilon)^2 < \lfloor M^2 \rfloor + 1$. We will build a function $D \in C_c(G \times X)$ such that $\text{Ind } \epsilon_z(D^* * D)$ is a rank-one projection and

$$\text{tr}(\text{Ind } \epsilon_{x_n}(D^* * D)) < M^2(1+\epsilon)^2 < \lfloor M^2 \rfloor + 1$$

frequently. (The function D is similar to the ones used in [20, Proposition 4.2] and [13, Proposition 4.5].) By the generalised lower semi-continuity result of [6, Theorem 4.3] we

will have

$$\begin{aligned} \liminf_n \operatorname{tr}(\operatorname{Ind} \epsilon_{x_n}(D^* * D)) &\geq M_L(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})) \operatorname{tr}(\operatorname{Ind} \epsilon_z(D^* * D)) \\ &= M_L(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})), \end{aligned}$$

and the theorem will follow.

Let $\delta > 0$ such that

$$\delta < \frac{\epsilon \nu(\phi_z^{-1}(V))}{1 + \epsilon} < \nu(\phi_z^{-1}(V)).$$

By the regularity of the measure ν there exists a compact subset W of the open set $\phi_z^{-1}(V)$ such that

$$0 < \nu(\phi_z^{-1}(V)) - \delta < \nu(W).$$

Since W is compact, there is a compact neighbourhood W_1 of W contained in $\phi_z^{-1}(V)$ and a continuous function $g : G \rightarrow [0, 1]$ such that g is identically one on W and is identically zero off the interior of W_1 . Then

$$\nu(\phi_z^{-1}(V)) - \delta < \nu(W) \leq \int_G g(t)^2 dt = \|g\|_2^2,$$

and hence

$$\frac{\nu(\phi_z^{-1}(V))}{\|g\|_2^2} < 1 + \frac{\delta}{\|g\|_2^2} < 1 + \frac{\delta}{\nu(\phi_z^{-1}(V)) - \delta} < 1 + \epsilon. \quad (3.1)$$

Since $G \cdot z$ is locally closed in X it follows from [9, Theorem 1], applied to the locally compact Hausdorff transformation group $(G, G \cdot z)$, that ϕ_z is a homeomorphism of G onto $G \cdot z$. So there is a continuous function $g_1 : W_1 \cdot z \rightarrow [0, 1]$ such that $g_1(t \cdot z) = g(t)$ for $t \in W_1$. Since $W_1 \cdot z$ is a compact subset of the locally compact Hausdorff space X , it follows from Tietze's Extension Theorem (applied to the one-point compactification of X if necessary) that g_1 can be extended to a continuous function $g_2 : X \rightarrow [0, 1]$. Because $W_1 \cdot z$ is a compact subset of the open set V , there exists a compact neighbourhood P of $W_1 \cdot z$ contained in V and a continuous function $h : X \rightarrow [0, 1]$ such that h is identically one on $W_1 \cdot z$ and is identically zero off the interior of P . Note that h has compact support contained in P . We set

$$f(x) = h(x)g_2(x).$$

Then $f \in C_c(X)$ with $0 \leq f \leq 1$ and $\operatorname{supp} f \subset \operatorname{supp} h \subset P \subset V$. Note that

$$\|f_z\|_2^2 = \int_G f(t \cdot z)^2 dt = \int_G h(t \cdot z)^2 g_2(t \cdot z)^2 dt \geq \int_{W_1} g(t)^2 dt = \|g\|_2^2 \quad (3.2)$$

since h is identically one on $W_1 \cdot z$ and g has support inside W_1 . We now set

$$F(x) = \frac{f(x)}{\|f_z\|_2}.$$

Now $F \in C_c(X)$, $\|F_z\|_2 = 1$ and $F_x(s) = F(s \cdot x) \neq 0$ implies $s \in \phi_x^{-1}(V)$ by our choice of h . Since $\phi_z^{-1}(V)$ is relatively compact, $\operatorname{supp} F_z$ is compact.

Choose $b \in C_c(G \times X)$ such that $0 \leq b \leq 1$ and b is identically one on the set $(\operatorname{supp} F_z)(\operatorname{supp} F_z)^{-1} \times \operatorname{supp} F$. Set

$$B(t, x) = F(x)F(t^{-1} \cdot x)b(t^{-1}, x)\Delta(t)^{-1/2} \quad \text{and} \quad D = \frac{1}{2}(B + B^*).$$

Using (2.1), we have

$$(\text{Ind } \epsilon_x(B)\xi)(s) = \int_G F(s \cdot x) F(w \cdot x) b(ws^{-1}, s \cdot x) \xi(w) d\nu(w)$$

for $\xi \in L^2(G, \nu)$. So

$$(\text{Ind } \epsilon_x(D)\xi)(s) = \frac{1}{2} F(s \cdot x) \int_G F(w \cdot x) (b(ws^{-1}, s \cdot x) + b(sw^{-1}, w \cdot x)) \xi(w) d\nu(w).$$

If $s, w \in \text{supp } F_z$ then $s \cdot z, w \cdot z \in \text{supp } F$ by the continuity of the action, and hence $b(ws^{-1}, s \cdot z) + b(sw^{-1}, w \cdot z) = 2$. It follows that $\text{Ind } \epsilon_z(D)(\xi) = (\xi, F_z) F_z$. Thus $\text{Ind } \epsilon_z(D)$, and hence $\text{Ind } \epsilon_z(D^* * D)$, is the rank-one projection determined by the unit vector $F_z \in L^2(G, \nu)$.

Choose a subsequence $(x_{n_i})_i$ of $(x_n)_n$ such that

$$\nu(\phi_{x_{n_i}}^{-1}(V)) \leq M\nu(\phi_z^{-1}(V))$$

for all $i \geq 1$ and set $E_i = \{s \in G : F(s \cdot x_{n_i}) \neq 0\}$. Then each E_i is open (hence measurable) with

$$\nu(E_i) \leq \nu(\phi_{x_{n_i}}^{-1}(V)) \leq M\nu(\phi_z^{-1}(V)) < \infty$$

and

$$\int_G F(s \cdot x_{n_i})^2 d\nu(s) \leq \frac{\nu(E_i)}{\|f_z\|_2^2} \leq \frac{M\nu(\phi_z^{-1}(V))}{\|g\|_2^2} \quad (3.3)$$

using (3.2). The function

$$S_i(s, w) := \frac{1}{2} F(s \cdot x_{n_i}) F(w \cdot x_{n_i}) (b(ws^{-1}, s \cdot x_{n_i}) + b(sw^{-1}, w \cdot x_{n_i}))$$

is continuous and is bounded by $\|F\|_\infty^2$ because $0 \leq b \leq 1$. Thus $S_i \in L^2(G \times G)$ and $\text{Ind } \epsilon_{x_{n_i}}(D)$ is the self-adjoint Hilbert-Schmidt operator on $L^2(G, \nu)$ with kernel S_i . It follows that $\text{Ind } \epsilon_{x_{n_i}}(D^* * D)$ is a trace-class operator with

$$\text{tr}(\text{Ind } \epsilon_{x_{n_i}}(D^* * D)) = \|S_i\|_2^2$$

(see, for example, [17, Proposition 3.4.16]). An application of Fubini's Theorem gives

$$\begin{aligned} & \text{tr}(\text{Ind } \epsilon_{x_{n_i}}(D^* * D)) \\ &= \frac{1}{4} \int \int F(s \cdot x_{n_i})^2 F(w \cdot x_{n_i})^2 (b(ws^{-1}, s \cdot x_{n_i}) + b(sw^{-1}, w \cdot x_{n_i}))^2 d\nu(w) d\nu(s) \\ &\leq \left(\int_G F(s \cdot x_{n_i})^2 d\nu(s) \right)^2 \\ &\leq \frac{M^2 \nu(\phi_z^{-1}(V))^2}{\|g\|_2^4} \quad (\text{using (3.3)}) \\ &< M^2(1 + \epsilon)^2 \quad (\text{using (3.1)}). \end{aligned} \quad (3.4)$$

Now

$$M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \leq \liminf_n \text{tr}(\text{Ind } \epsilon_{x_n}(D^* * D)) \leq M^2(1 + \epsilon)^2 < \lfloor M^2 \rfloor + 1,$$

and hence $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \leq \lfloor M^2 \rfloor$. □

Our aim is to use Theorem 3.1 to bootstrap to a result where the relative lower multiplicity is bounded by $\lfloor M \rfloor$ rather than $\lfloor M^2 \rfloor$ (see Theorem 3.5). To do this we will need to be able to cut away any limits of $(\text{Ind } \epsilon_{x_n})_n$ that are different from $\text{Ind } \epsilon_z$ so that we can apply the following crucial Lemma 3.2.

Lemma 3.2. *Suppose that (G, X) is a transformation group. Let W be a compact neighbourhood of $z \in X$, K a compact subset of G , and U an open neighbourhood of $\phi_z^{-1}(W)$ in G . Let $(x_n)_{n \geq 1}$ be a sequence in X such that $G \cdot x_n \rightarrow G \cdot z$ and $G \cdot z$ is the unique limit of $(G \cdot x_n)_{n \geq 1}$ in X/G . There exists n_0 such that for every $n \geq n_0$ and every $s \in \phi_{x_n}^{-1}(W)$ there exists $r \in \phi_z^{-1}(W)$ such that $Ks \cap \phi_{x_n}^{-1}(W) \subset Ur^{-1}s$.*

Proof. Suppose that there is no such n_0 . Then, by passing to a subsequence, for each n there exists $s_n \in \phi_{x_n}^{-1}(W)$ such that

$$Ks_n \cap \phi_{x_n}^{-1}(W) \not\subset Ur^{-1}s_n \quad (3.5)$$

for all $r \in \phi_z^{-1}(W)$.

Note that $s_n \cdot x_n \in W$, so by passing to a further subsequence we may assume that $s_n \cdot x_n$ converges to some $y \in W$. Since $G \cdot z$ is the unique limit point of $(G \cdot x_n)$ in X/G , we have $y = r \cdot z$ for some $r \in G$. Since $y \in W$ we have $r \in \phi_z^{-1}(W)$, and for this r Equation 3.5 holds for all n . So there exist $k_n \in K$ such that $k_n s_n \in \phi_{x_n}^{-1}(W)$ but $k_n \notin Ur^{-1}$. By passing to a further subsequence $(k_n)_n$ converges to $k \in K$. Then $k_n s_n \cdot x_n \rightarrow k \cdot y = kr \cdot z \in W$. But now $k \in \phi_z^{-1}(W)r^{-1} \subset Ur^{-1}$ and $k_n \notin Ur^{-1}$ which is impossible since U is open and $k_n \rightarrow k$. $\square$

In the next Lemma we consider a potentially bad neighbourhood V of z , where the measure of $\phi_z^{-1}(\overline{V})$ might be much larger than that of $\phi_z^{-1}(V)$, and show that we can at least find a nicer, controlled, neighbourhood contained in it.

Lemma 3.3. *Suppose that (G, X) is a transformation group. Let $\gamma > 0$, $z \in Z$ and let V be an open neighbourhood of z in X such that $\nu(\phi_z^{-1}(V)) < \infty$. Then there exists an open relatively compact neighbourhood V_1 of z such that $\overline{V_1} \subset V$ and*

$$\nu(\phi_z^{-1}(V)) - \gamma < \nu(\phi_z^{-1}(V_1)) \leq \nu(\phi_z^{-1}(\overline{V_1})) \leq \nu(\phi_z^{-1}(V)) < \nu(\phi_z^{-1}(V_1)) + \gamma. \quad (3.6)$$

Proof. By the regularity of the measure ν we can choose a compact subset W of $\phi_z^{-1}(V)$ such that $e \in W$ and $\nu(W) > \nu(\phi_z^{-1}(V)) - \gamma$. Note that $W \cdot z$ is a compact subset of $G \cdot z$ and hence of X . Hence there exists an open relatively compact neighbourhood V_1 of $W \cdot z$ such that $\overline{V_1} \subset V$. Now

$$\nu(\phi_z^{-1}(V)) - \gamma < \nu(W) \leq \nu(\phi_z^{-1}(V_1)) \leq \nu(\phi_z^{-1}(\overline{V_1})) \leq \nu(\phi_z^{-1}(V))$$

and (3.6) follows. $\square$

The following result is a sharpening of [5, Proposition 2.1(i)]. We will use it to cut away multiple limits of our sequences.

Proposition 3.4. *Let A be a C^* -algebra and let $\pi \in \hat{A}$. Suppose that $\Omega = (\pi_\alpha)_{\alpha \in \Lambda}$ is a net in $\hat{A}$ which is convergent to π and that $M_L(\pi, \Omega) = k < \infty$. Then $\{\pi\}$ is open in the set $L(\Omega)$ of limits of Ω .*

Proof. The direct way to see this is to re-work the proof of [5, Proposition 2.1(i)]. An alternative shorter argument is as follows. By [6, Proposition 2.3], there is a subnet Ω_1 of Ω such that

$$M_U(\pi, \Omega_1) = M_L(\pi, \Omega) < \infty.$$

By [5, Proposition 2.1(i)], $\{\pi\}$ is open in $L(\Omega_1)$. But $L(\Omega) \subset L(\Omega_1)$ and so $\{\pi\}$ is open in $L(\Omega)$. $\square$

Theorem 3.5 has the same hypotheses as Theorem 3.1 but a stronger conclusion; the proof requires Theorem 3.1 and some estimates based on Lemmas 3.2 and 3.3.

Theorem 3.5. *Suppose that (G, X) is a free second-countable transformation group. Let $M \in \mathbb{R}$ with $M \geq 1$, let $z \in X$ and let $(x_n)_{n \geq 1}$ be a sequence in X . Assume that $G \cdot z$ is locally closed in X . Suppose that there exists an open neighbourhood V of z in X such that $\phi_z^{-1}(V)$ is relatively compact and*

$$\nu(\phi_{x_n}^{-1}(V)) \leq M\nu(\phi_z^{-1}(V))$$

frequently. Then $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \leq \lfloor M \rfloor$.

Proof. If $\text{Ind } \epsilon_{x_n} \not\rightarrow \text{Ind } \epsilon_z$ then $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) = 0 < \lfloor M \rfloor$. So we assume from now on that $\text{Ind } \epsilon_{x_n} \rightarrow \text{Ind } \epsilon_z$.

Our first claim is that $G \cdot x_n \rightarrow G \cdot z$. To see this, suppose that $G \cdot x_n \not\rightarrow G \cdot z$. Then there exists an open neighbourhood U_0 of $G \cdot z$ such that $G \cdot x_n$ is frequently not in U_0 . Let $q : X \rightarrow X/G$ be the quotient map. Then $U_1 = q^{-1}(U_0)$ is an open G -invariant neighbourhood of z and $x_n \notin U_1$ frequently. Note that $C_0(U_1) \times G$ is isomorphic to a closed two-sided ideal I of $C_0(X) \times G$ and $I \subset \ker(\text{Ind } \epsilon_{x_n})$ whenever $x_n \notin U_1$. Hence $\text{Ind } \epsilon_{x_n} \notin \hat{I}$ frequently. But $\hat{I}$ is an open neighbourhood of $\text{Ind } \epsilon_z$, so $\text{Ind } \epsilon_{x_n} \not\rightarrow \text{Ind } \epsilon_z$.

Next we claim that we may assume, without loss of generality, that $G \cdot z$ is the unique limit of $(G \cdot x_n)_n$ in X/G . To see this, note that $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \leq \lfloor M^2 \rfloor < \infty$ by Theorem 3.1. Hence, by Proposition 3.4, $\{\text{Ind } \epsilon_z\}$ is open in the set of limits of $(\text{Ind } \epsilon_{x_n})_n$. So there is an open neighbourhood U_2 of $\text{Ind } \epsilon_z$ in the spectrum of $C_0(X) \times G$ such that $\text{Ind } \epsilon_z$ is the unique limit of $(\text{Ind } \epsilon_{x_n})_n$ in U_2 .

Write $(X/G)^\sim$ for the T_0 -isation of X/G , so that $(X/G)^\sim$ is the quotient of X obtained by identifying points of X with equal orbit closures. It follows, for example from [19, Lemmas 4.5 and 4.10] that the map

$$(X/G)^\sim \rightarrow \text{Prim}(C_0(X) \times G)$$

given by $[x] \mapsto \ker(\text{Ind } \epsilon_x)$ is a homeomorphism. Since the topology on $(C_0(X) \times G)^\wedge$ is the topology pulled back from $\text{Prim}(C_0(X) \times G)$, it is straightforward to see that $\text{Ind} : X \rightarrow (C_0(X) \times G)^\wedge : x \mapsto \text{Ind } \epsilon_x$ is continuous. Let

$$Y = \text{Ind}^{-1}(U_2).$$

Then Y is an open G -invariant neighbourhood of z in X . Note that $x_n \in Y$ eventually.

Now suppose that, for some $y \in Y$, $G \cdot x_n \rightarrow G \cdot y$ in Y/G and hence in X/G . Then $\text{Ind } \epsilon_{x_n} \rightarrow \text{Ind } \epsilon_y$, and $\text{Ind } \epsilon_y \in U_2$ since $y \in \text{Ind}^{-1}(U_2)$. But $(\text{Ind } \epsilon_{x_n})$ has the unique limit $\text{Ind } \epsilon_z$ in U_2 , so $\text{Ind } \epsilon_z = \text{Ind } \epsilon_y$ and hence $\overline{G \cdot z} = \overline{G \cdot y}$ in X . Since $G \cdot z$ is locally closed we obtain $G \cdot z = G \cdot y$ in X and hence in Y . (To see this, note that there exists a sequence $(r_k)_k \subset G$ such that $r_k \cdot y \rightarrow z$. Let U_3 be an open subset of X such that $G \cdot z = U_3 \cap \overline{G \cdot z}$. Then $r_k \cdot y \in U_3$ eventually, so eventually $r_k \cdot y \in U_3 \cap \overline{G \cdot y} = U_3 \cap \overline{G \cdot z} = G \cdot z$.)

Finally, we note that $C_0(Y) \times G$ is isomorphic to a closed two-sided ideal J of $C_0(X) \times G$ and $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n}))$ is the same whether we compute it in the ideal J or in $C_0(X) \times G$ (see [5, Proposition 5.3]); moreover, $\phi_z^{-1}(V) = \phi_z^{-1}(V \cap Y)$ and $\phi_{x_n}^{-1}(V) = \phi_{x_n}^{-1}(V \cap Y)$ when n is large enough so that $x_n \in Y$. Thus we may replace X by Y and therefore assume that $G \cdot z$ is the unique limit of $G \cdot x_n$ in X/G , as claimed.

The idea of the rest of the proof is the same as in Theorem 3.1, but our estimates are more delicate. Fix $\epsilon > 0$ such that $M(1 + \epsilon)^2 < \lfloor M \rfloor + 1$ and choose $\gamma > 0$ such that

$$\gamma < \frac{\epsilon \nu(\phi_z^{-1}(V))}{1 + \epsilon} < \nu(\phi_z^{-1}(V)). \quad (3.7)$$

By Lemma 3.3 there exists an open relatively compact neighbourhood V_1 of z such that $\overline{V_1} \subset V$ and

$$0 < \nu(\phi_z^{-1}(V)) - \gamma < \nu(\phi_z^{-1}(V_1)) \leq \nu(\phi_z^{-1}(\overline{V_1})) \leq \nu(\phi_z^{-1}(V)) < \nu(\phi_z^{-1}(V_1)) + \gamma.$$

(The reason for passing from V to V_1 is that we will later apply Lemma 3.2 to the compact neighbourhood $\overline{V_1}$ and, in contrast to what could happen with $\overline{V}$, we can control $\nu(\phi_z^{-1}(\overline{V_1}))$ relative to $\nu(\phi_z^{-1}(V_1))$.) Choose a subsequence $(x_{n_i})_i$ of $(x_n)_n$ such that

$$\nu(\phi_{x_{n_i}}^{-1}(V)) \leq M \nu(\phi_z^{-1}(V))$$

for all $i \geq 1$. Then

$$\begin{aligned} \nu(\phi_{x_{n_i}}^{-1}(V_1)) &\leq \nu(\phi_{x_{n_i}}^{-1}(V)) \\ &\leq M \nu(\phi_z^{-1}(V)) \quad (\text{by assumption}) \\ &< M(\nu(\phi_z^{-1}(V_1)) + \gamma) \\ &< M \nu(\phi_z^{-1}(V_1)) + M \epsilon (\nu(\phi_z^{-1}(V)) - \gamma) \quad (\text{by (3.7)}) \\ &< M \nu(\phi_z^{-1}(V_1)) + M \epsilon \nu(\phi_z^{-1}(V_1)) \\ &= M(1 + \epsilon) \nu(\phi_z^{-1}(V_1)) \end{aligned} \quad (3.8)$$

for all i . Since

$$\frac{\nu(\phi_z^{-1}(V_1))(\nu(\phi_z^{-1}(V_1)) + \gamma + \frac{1}{j})}{(\nu(\phi_z^{-1}(V_1)) - \frac{1}{j})^2} \rightarrow 1 + \frac{\gamma}{\nu(\phi_z^{-1}(V_1))} < 1 + \epsilon$$

as $j \rightarrow \infty$, there exists $\delta > 0$ such that $\delta < \nu(\phi_z^{-1}(V_1))$ and

$$\frac{\nu(\phi_z^{-1}(V_1))(\nu(\phi_z^{-1}(\overline{V_1})) + \delta)}{(\nu(\phi_z^{-1}(V_1)) - \delta)^2} < \frac{\nu(\phi_z^{-1}(V_1))(\nu(\phi_z^{-1}(V_1)) + \gamma + \delta)}{(\nu(\phi_z^{-1}(V_1)) - \delta)^2} < 1 + \epsilon. \quad (3.9)$$

Next we construct a function $F \in C_c(X)$ with support inside V_1 . By the regularity of the measure ν there exists a compact subset W of the open set $\phi_z^{-1}(V_1)$ such that $0 < \nu(\phi_z^{-1}(V_1)) - \delta < \nu(W)$. Since W is compact, there is a compact neighbourhood W_1 of W contained in $\phi_z^{-1}(V_1)$ and a continuous function $g : G \rightarrow [0, 1]$ such that g is identically one on W and is identically zero off the interior of W_1 . Then

$$\nu(\phi_z^{-1}(V_1)) - \delta < \nu(W) \leq \int_G g(t)^2 dt = \|g\|_2^2. \quad (3.10)$$

There is a continuous function $g_1 : W_1 \cdot z \rightarrow [0, 1]$ such that $g_1(t \cdot z) = g(t)$ for $t \in W_1$. Since $W_1 \cdot z$ is a compact subset of X , we can extend g_1 to a continuous function $g_2 : X \rightarrow [0, 1]$

using Tietze's Extension Theorem. There exists a compact neighbourhood P of $W_1 \cdot z$ contained in V_1 and a continuous function $h : X \rightarrow [0, 1]$ such that h is identically one on $W_1 \cdot z$ and is identically zero off the interior of P . Note that h has compact support contained in P . We set $f(x) = h(x)g_2(x)$. Then f is continuous on X and

$$\|f_z\|_2^2 = \int_G f(t \cdot z)^2 dt = \int_G h(t \cdot z)^2 g_2(t \cdot z)^2 dt \geq \int_{W_1} g(t)^2 dt = \|g\|_2^2 \quad (3.11)$$

since h is identically one on $W_1 \cdot z$ and g has support inside W_1 . We now set $F(x) = \frac{f(x)}{\|f_z\|_2}$. Thus $F \in C_c(X)$, $\|F_z\|_2 = 1$ and $F_x(s) = F(s \cdot x) \neq 0$ implies $s \in \phi_x^{-1}(V_1)$ by our choice of h .

Let K be an open relatively compact symmetric neighbourhood of $(\text{supp } F_z)(\text{supp } F_z)^{-1}$ in G and L an open relatively compact neighbourhood of $\text{supp } F$ in X . Choose $b \in C_c(G \times X)$ such that $0 \leq b \leq 1$ and b is identically one on the set $(\text{supp } F_z)(\text{supp } F_z)^{-1} \times \text{supp } F$ and b is identically zero off $K \times L$. (Thus b is as in Theorem 3.1, but we have rounded it off.) Set

$$B(t, x) = F(x)F(t^{-1} \cdot x)b(t^{-1}, x)\Delta(t)^{-1/2} \quad \text{and} \quad D = \frac{1}{2}(B + B^*).$$

Again, $\text{Ind } \epsilon_z(D)$, and hence $\text{Ind } \epsilon_z(D^* * D)$, is the rank-one projection determined by the unit vector $F_z \in L^2(G, \nu)$. From (3.4) we have

$$\begin{aligned} & \text{tr}(\text{Ind } \epsilon_{x_{n_i}}(D^* * D)) \\ &= \frac{1}{4} \int_G F(s \cdot x_{n_i})^2 \left(\int_G F(w \cdot x_{n_i})^2 (b(ws^{-1}, s \cdot x_{n_i}) + b(sw^{-1}, w \cdot x_{n_i}))^2 d\nu(w) \right) d\nu(s). \end{aligned}$$

The inner integrand is zero unless $w \in \phi_{x_{n_i}}^{-1}(V_1) \cap Ks$ because $F_{x_{n_i}}(w) = F(w \cdot x_{n_i}) \neq 0$ implies $w \in \phi_{x_{n_i}}^{-1}(V_1)$ and b is identically zero off $K \times L$. Thus

$$\begin{aligned} \text{tr}(\text{Ind } \epsilon_{x_{n_i}}(D^* * D)) &\leq \int_{s \in \phi_{x_{n_i}}^{-1}(V_1)} F(s \cdot x_{n_i})^2 \left(\int_{w \in \phi_{x_{n_i}}^{-1}(V_1) \cap Ks} F(w \cdot x_{n_i})^2 d\nu(w) \right) d\nu(s) \\ &\leq \frac{1}{\|f_z\|_2^4} \int_{s \in \phi_{x_{n_i}}^{-1}(V_1)} 1 \left(\int_{w \in \phi_{x_{n_i}}^{-1}(V_1) \cap Ks} 1 d\nu(w) \right) d\nu(s). \end{aligned}$$

Choose an open neighbourhood U of $\phi_z^{-1}(\overline{V_1})$ such that $\nu(U) < \nu(\phi_z^{-1}(\overline{V_1})) + \delta$. By Lemma 3.2, applied with $\overline{V_1}$, $\overline{K}$ and U , there exists i_0 such that, for every $i \geq i_0$ and every $s \in \phi_{x_{n_i}}^{-1}(\overline{V_1})$ there exists $r \in \phi_z^{-1}(\overline{V_1})$ with $\overline{K}s \cap \phi_{x_{n_i}}^{-1}(\overline{V_1}) \subset Ur^{-1}s$. It follows that

$$\nu(\overline{K}s \cap \phi_{x_{n_i}}^{-1}(\overline{V_1})) \leq \nu(U) < \nu(\phi_z^{-1}(\overline{V_1})) + \delta$$

by the right-invariance of ν . So, provided $i \geq i_0$,

$$\begin{aligned}
\operatorname{tr}(\operatorname{Ind} \epsilon_{x_{n_i}}(D^* * D)) &\leq \frac{\nu(\phi_{x_{n_i}}^{-1}(V_1))(\nu(\phi_z^{-1}(\overline{V_1})) + \delta)}{\|f_z\|_2^4} \\
&< \frac{M(1 + \epsilon)\nu(\phi_z^{-1}(V_1))(\nu(\phi_z^{-1}(\overline{V_1})) + \delta)}{\|g\|_2^4} && \text{using (3.8) and (3.11)} \\
&\leq \frac{M(1 + \epsilon)\nu(\phi_z^{-1}(V_1))(\nu(\phi_z^{-1}(\overline{V_1})) + \delta)}{(\nu(\phi_z^{-1}(V_1)) - \delta)^2} && \text{using (3.10)} \\
&< M(1 + \epsilon)^2 && \text{using (3.9).}
\end{aligned}$$

By generalised lower semi-continuity [6, Theorem 4.3]

$$\begin{aligned}
\liminf_n \operatorname{tr}(\operatorname{Ind} \epsilon_{x_n}(D^* * D)) &\geq M_L(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})) \operatorname{tr}(\operatorname{Ind} \epsilon_z(D^* * D)) \\
&= M_L(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})).
\end{aligned}$$

We now have

$$M_L(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})) \leq \liminf_n \operatorname{tr}(\operatorname{Ind} \epsilon_{x_n}(D^* * D)) \leq M(1 + \epsilon)^2 < \lfloor M \rfloor + 1,$$

and hence $M_L(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})) \leq \lfloor M \rfloor$. $\square$

The first corollary of Theorem 3.5 is the analogous result for the upper multiplicity relative to a net.

Theorem 3.6. *Suppose that (G, X) is a free second-countable transformation group. Let $M \in \mathbb{R}$ with $M \geq 1$, let $z \in X$ and let $(x_n)_{n \geq 1}$ be a sequence in X . Assume that $G \cdot z$ is locally closed in X . Suppose that there exists an open neighbourhood V of z in X such that $\phi_z^{-1}(V)$ is relatively compact and*

$$\nu(\phi_{x_n}^{-1}(V)) \leq M\nu(\phi_z^{-1}(V)) < \infty$$

eventually. Then $M_U(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})) \leq \lfloor M \rfloor$.

Proof. Both G and X are second countable, so $C_0(X) \times G$ is separable. By Lemma A.1 there exists a subsequence $(\operatorname{Ind} \epsilon_{x_{n_i}})_i$ such that

$$M_U(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})) = M_U(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_{n_i}})) = M_L(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_{n_i}})).$$

By Theorem 3.5, $M_L(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_{n_i}})) \leq \lfloor M \rfloor$, and hence $M_U(\operatorname{Ind} \epsilon_z, (\operatorname{Ind} \epsilon_{x_n})) \leq \lfloor M \rfloor$. $\square$

Corollary 3.7. *Suppose that (G, X) is a free second-countable transformation group such that all the orbits are locally closed in X . Let $M \in \mathbb{R}$ with $M \geq 1$ and let $z \in X$. If for every sequence $(x_n)_{n \geq 1}$ in X which converges to z there exists an open neighbourhood V of z in X such that $\phi_z^{-1}(V)$ is relatively compact and*

$$\nu(\phi_{x_n}^{-1}(V)) \leq M\nu(\phi_z^{-1}(V)) < \infty$$

frequently, then $M_U(\operatorname{Ind} \epsilon_z) \leq \lfloor M \rfloor$.

Proof. Since $C_0(X) \times G$ is separable it follows from [3, Lemma 1.2] that there exists a sequence $(\pi_n)_{n \geq 1}$ in $(C_0(X) \times G)^\wedge$ converging to $\text{Ind } \epsilon_z$, such that

$$M_L(\text{Ind } \epsilon_z, (\pi_n)) = M_U(\text{Ind } \epsilon_z, (\pi_n)) = M_U(\text{Ind } \epsilon_z).$$

Since the orbits are locally closed, $C_0(X) \times G$ is Type I by [10, Theorem 3.3], and then the map $x \mapsto \text{Ind } \epsilon_x$ induces a homeomorphism of X/G onto $C_0(X) \times G)^\wedge$ by [11, Lemma 16]. In particular, the mapping $X \rightarrow (C_0(X) \times G)^\wedge$ is an open surjection so there exists a sequence $(x_i)_{i \geq 1}$ in X converging to z such that $(\text{Ind } \epsilon_{x_i})_{i \geq 1}$ is a subsequence of $(\pi_n)_{n \geq 1}$. By Theorem 3.5, $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_i})) \leq \lfloor M \rfloor$. Since

$$\begin{aligned} M_U(\text{Ind } \epsilon_z) &= M_L(\text{Ind } \epsilon_z, (\pi_n)) \leq M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_i})) \leq M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_i})) \\ &\leq M_U(\text{Ind } \epsilon_z, (\pi_n)) = M_U(\text{Ind } \epsilon_z), \end{aligned}$$

we obtain $M_U(\text{Ind } \epsilon_z) \leq \lfloor M \rfloor$. $\square$

4. TOPOLOGICAL STRENGTH OF CONVERGENCE

A sequence $(t_n)_{n \geq 1} \subset G$ tends to infinity if it admits no convergent subsequence. Let $k \in \mathbb{P}$. A sequence $(x_n)_{n \geq 1}$ in X is *k-times convergent in X/G to $z \in X$* if there exist k sequences $(t_n^{(1)})_n, (t_n^{(2)})_n, \dots, (t_n^{(k)})_n \subset G$, such that

- (1) $t_n^{(i)} \cdot x_n \rightarrow z$ as $n \rightarrow \infty$ for $1 \leq i \leq k$, and
- (2) if $1 \leq i < j \leq k$ then $t_n^{(j)}(t_n^{(i)})^{-1} \rightarrow \infty$ as $n \rightarrow \infty$.

This definition of *k-times convergence* is a special case of [2, Definition 2.2] obtained by taking $A = C_0(X)$ and replacing $t_n^{(i)}$ by $(t_n^{(i)})^{-1}$.

Proposition 4.1 below shows that measure-theoretic accumulation implies topological strength of convergence in X/G ; the proof uses our key technical Lemma 3.2.

Proposition 4.1. *Let (G, X) be a second-countable transformation group. Let $k \in \mathbb{P}$ and $z \in X$, with $G \cdot z$ locally closed in X . Assume that $(x_n)_{n \geq 1}$ is a sequence in X such that $G \cdot x_n \rightarrow G \cdot z$ and $G \cdot z$ is the unique limit of $(G \cdot x_n)_n$ in X/G .*

- (1) *Suppose that there exists a basic sequence $(W_m)_{m \geq 1}$ of compact neighbourhoods of z with $W_{m+1} \subset W_m$, such that, for each m*

$$\liminf_n \nu(\phi_{x_n}^{-1}(W_m)) > (k-1)\nu(\phi_z^{-1}(W_m)).$$

Then (x_n) converges k-times in X/G to z .

- (2) *Suppose that there exists a basic sequence $(W_m)_{m \geq 1}$ of compact neighbourhoods of z with $W_{m+1} \subset W_m$, such that, for each m*

$$\limsup_n \nu(\phi_{x_n}^{-1}(W_m)) > (k-1)\nu(\phi_z^{-1}(W_m)).$$

Then there exists a subsequence of $(x_n)_n$ which converges k-times in X/G to z .

Proof. Let $(K_m)_{m \geq 1}$ be an increasing sequence of compact subsets of G such that $G = \bigcup_{m \geq 1} \text{Int}(K_m)$.

- (1) By regularity of ν , for each $m \geq 1$, there exists an open neighbourhood U_m of $\phi_z^{-1}(W_m)$ such that

$$\liminf_n \nu(\phi_{x_n}^{-1}(W_m)) > (k-1)\nu(U_m). \quad (4.1)$$

We will construct, by induction, a strictly increasing sequence of positive integers $(n_m)_{m \geq 1}$ such that, for all $n \geq n_m$,

$$\nu(K_m s \cap \phi_{x_n}^{-1}(W_m)) \leq \nu(U_m) \quad \text{for all } s \in \phi_{x_n}^{-1}(W_m), \text{ and} \quad (4.2)$$

$$\nu(\phi_{x_n}^{-1}(W_m)) > (k-1)\nu(U_m). \quad (4.3)$$

We construct n_1 by applying Lemma 3.2 to K_1, W_1 and U_1 to obtain (4.2) for $m = 1$, and then, if necessary, increasing n_1 to ensure that (4.3) holds using (4.1) with $m = 1$. Assuming that we have constructed $n_1 < n_2 < \dots < n_{m-1}$, we apply Lemma 3.2 to K_m, W_m and U_m to obtain $n_m > n_{m-1}$ such that (4.2) holds, and again, if necessary, increase n_m to obtain (4.3).

If $n_1 > 1$ then, for $1 \leq n < n_1$, we set $t_n^{(i)} = e$ for $1 \leq i \leq k$. For each $n \geq n_1$ there is a unique m such that $n_m \leq n < n_{m+1}$. Choose $t_n^{(1)} \in \phi_{x_n}^{-1}(W_m)$. Using (4.2) and (4.3)

$$\nu(\phi_{x_n}^{-1}(W_m) \setminus K_m t_n^{(1)}) > (k-2)\nu(U_m).$$

So we may choose $t_n^{(2)} \in \phi_{x_n}^{-1}(W_m) \setminus K_m t_n^{(1)}$. Continuing in this way, we use (4.2) and (4.3) to choose

$$t_n^{(3)} \in \phi_{x_n}^{-1}(W_m) \setminus (K_m t_n^{(1)} \cup K_m t_n^{(2)})$$

$\vdots$

$$t_n^{(k)} \in \phi_{x_n}^{-1}(W_m) \setminus (\cup_{i=1}^{k-1} K_m t_n^{(i)}).$$

Note that for $n_m \leq n < n_{m+1}$ we have

$$t_n^{(i)} \cdot x_n \in W_m \text{ for } 1 \leq i \leq k \quad \text{and} \quad t_n^{(j)} \notin K_m t_n^{(i)} \text{ for } 1 \leq i < j \leq k.$$

We claim that $t_n^{(i)} \cdot x_n \rightarrow z$ as $n \rightarrow \infty$ for $1 \leq i \leq k$. To see this, fix i and let V be a neighbourhood of z . There exists m_0 such that $W_m \subset V$ for all $m \geq m_0$. For each $n \geq n_{m_0}$ there exists a $m \geq m_0$ such that $n_m \leq n < n_{m+1}$, and thus $t_n^{(i)} \cdot x_n \in W_m \subset V$.

Next, we claim that $t_n^{(j)}(t_n^{(i)})^{-1} \rightarrow \infty$ as $n \rightarrow \infty$ for $1 \leq i < j \leq k$. Fix $i < j$ and let K be a compact subset of G . There exists m_0 such that $K_m \supset K$ for all $m \geq m_0$. Then for $n \geq n_{m_0}$ there exists a $m \geq m_0$ such that $n_m \leq n < n_{m+1}$, and hence $t_n^{(j)}(t_n^{(i)})^{-1} \in G \setminus K_m \subset G \setminus K$. We have shown that $(x_n)_n$ converges k -times in X/G to z .

(2) By regularity of ν , for each $m \geq 1$ there exists an open neighbourhood U_m of $\phi_z^{-1}(W_m)$ such that

$$\limsup_n \nu(\phi_{x_n}^{-1}(W_m)) > (k-1)\nu(U_m). \quad (4.4)$$

We will construct, by induction, an increasing sequence $(i_m)_{m \geq 1}$ such that

$$\nu(K_m s \cap \phi_{x_{i_m}}^{-1}(W_m)) \leq \nu(U_m) \text{ for all } s \in \phi_{x_{i_m}}^{-1}(W_m) \text{ and} \quad (4.5)$$

$$\nu(\phi_{x_{i_m}}^{-1}(W_m)) > (k-1)\nu(U_m). \quad (4.6)$$

To start, we apply Lemma 3.2 to K_1, W_1 and U_1 to obtain n_1 such that

$$n \geq n_1 \text{ implies } \nu(K_1 s \cap \phi_{x_n}^{-1}(W_1)) \leq \nu(U_1) \text{ for all } s \in \phi_{x_n}^{-1}(W_1).$$

Then, using (4.4) with $m = 1$, choose $i_1 \geq n_1$ such that

$$\nu(\phi_{x_{i_1}}^{-1}(W_1)) > (k-1)\nu(U_1).$$

Assuming that $i_1 < i_2 < \dots < i_{m-1}$ have been chosen, apply Lemma 3.2 to K_m, W_m and U_m to obtain $n_m > i_{m-1}$ such that

$$n \geq n_m \text{ implies } \nu(K_m s \cap \phi_{x_n}^{-1}(W_m)) \leq \nu(U_m) \text{ for all } s \in \phi_{x_n}^{-1}(W_m),$$

and then, using (4.4), choose $i_m \geq n_m$ such that

$$\nu(\phi_{x_{i_m}}^{-1}(W_m)) > (k-1)\nu(U_m).$$

Now, for each $m \geq 1$, choose $t_{i_m}^{(1)} \in \phi_{x_{i_m}}^{-1}(W_m)$. Using (4.5) and (4.6)

$$\nu(\phi_{x_{i_m}}^{-1}(W_m) \setminus K_m t_{i_m}^{(1)}) > (k-2)\nu(U_m).$$

So we may choose $t_{i_m}^{(2)} \in \phi_{x_{i_m}}^{-1}(W_m) \setminus K_m t_{i_m}^{(1)}$. Continuing in this way, we use (4.5) and (4.6) to choose

$$\begin{aligned} t_{i_m}^{(3)} &\in \phi_{x_{i_m}}^{-1}(W_m) \setminus (K_m t_{i_m}^{(1)} \cup K_m t_{i_m}^{(2)}) \\ &\vdots \\ t_{i_m}^{(k)} &\in \phi_{x_{i_m}}^{-1}(W_m) \setminus (\cup_{j=1}^{k-1} K_m t_{i_m}^{(j)}). \end{aligned}$$

Since $t_{i_m}^{(j)} \cdot x_{i_m} \in W_m$ and $(W_m)_m$ is decreasing, it follows that $t_{i_m}^{(j)} \cdot x_{i_m} \rightarrow_m z$ for $1 \leq j \leq k$. If K is any compact subset of G then there exists m_0 such that $K \subset K_m$ for all $m \geq m_0$. Then for $m \geq m_0$ we have $t_{i_m}^{(j)}(t_{i_m}^{(l)})^{-1} \in G \setminus K_m \subset G \setminus K$ for $1 \leq l < j \leq k$. Thus $(x_{i_m})_m$ is a subsequence of $(x_n)_n$ which converges k -times in X/G to z . $\square$

Although we shall improve Proposition 4.2 later (see Corollary 5.6), we prove it now because it will be needed in the proof of Theorem 1.1.

Proposition 4.2. *Suppose that (G, X) is a free second-countable transformation group. Let $z \in X$ and let $(x_n)_{n \geq 1}$ be a sequence in X . Assume that $G \cdot z$ is locally closed in X . Consider the following properties.*

- (1) $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) = \infty$;
- (2) *for every open neighbourhood V of z such that $\phi_z^{-1}(V)$ is relatively compact, $\nu(\phi_{x_n}^{-1}(V)) \rightarrow \infty$ as $n \rightarrow \infty$;*
- (3) *for each $k \geq 1$, the sequence $(x_n)_n$ converges k -times in X/G to z .*

Then (1) $\implies$ (2) and (2) $\implies$ (3).

Proof. (1) $\implies$ (2). This is an immediate corollary of Theorem 3.1.

(2) $\implies$ (3). Let $(K_m)_{m \geq 1}$ be an increasing sequence of compact subsets of G such that $G = \cup_{m \geq 1} \text{Int}(K_m)$ and let $(V_m)_{m \geq 1}$ be a decreasing sequence of open, basic neighbourhoods of z such that $\phi_z^{-1}(V_1)$ is relatively compact (such neighbourhoods exist by Lemma 2.1). Let $k \geq 1$. Assuming (2), we have

$$\nu(\phi_{x_n}^{-1}(V_m)) \rightarrow_n \infty \quad (m \geq 1).$$

Hence we can construct inductively a strictly increasing sequence of positive integers $(n_m)_{m \geq 1}$ such that, for all $n \geq n_m$,

$$\nu(\phi_{x_n}^{-1}(V_m)) > (k-1)\nu(K_m). \tag{4.7}$$

We can now choose $t_n^{(i)}$ for $1 \leq i \leq k$ as in Proposition 4.1(1) by using (4.7) in place of (4.2) and (4.3). $\square$

5. MAIN RESULTS

We will shortly combine the results from §3 and §4 to prove our main theorem stated in the introduction.

Lemma 5.1. *Suppose that (G, X) is a transformation group. Let $k \in \mathbb{P}$, $z \in X$ and $(x_n)_{n \geq 1}$ be a sequence in X . Assume that $G \cdot z$ is locally closed in X and that there exists a real number $R > k - 1$ such that for every open neighbourhood U of z with $\phi_z^{-1}(U)$ relatively compact we have*

$$\liminf_n \nu(\phi_{x_n}^{-1}(U)) \geq R\nu(\phi_z^{-1}(U)).$$

Given an open neighbourhood V of z such that $\phi_z^{-1}(V)$ is relatively compact, there exists a compact neighbourhood N of z with $N \subset V$ such that

$$\liminf_n \nu(\phi_{x_n}^{-1}(N)) > (k - 1)\nu(\phi_z^{-1}(N)).$$

Proof. Apply Lemma 3.3 to V with $0 < \gamma < \left(\frac{R-k+1}{R}\right)\nu(\phi_z^{-1}(V))$ to get an open relatively compact neighbourhood V_1 of z with $\overline{V_1} \subset V$ and

$$\nu(\phi_z^{-1}(V)) - \gamma < \nu(\phi_z^{-1}(V_1)) \leq \nu(\phi_z^{-1}(\overline{V_1})) \leq \nu(\phi_z^{-1}(V)) < \nu(\phi_z^{-1}(V_1)) + \gamma.$$

Since $\phi_z^{-1}(V_1)$ is relatively compact we have

$$\begin{aligned} \liminf_n \nu(\phi_{x_n}^{-1}(\overline{V_1})) &\geq \liminf_n \nu(\phi_{x_n}^{-1}(V_1)) \\ &\geq R\nu(\phi_z^{-1}(V_1)) \quad \text{by hypothesis} \\ &> R(\nu(\phi_z^{-1}(V)) - \gamma) \\ &> (k - 1)\nu(\phi_z^{-1}(V)) \quad \text{by our choice of } \gamma \\ &\geq (k - 1)\nu(\phi_z^{-1}(\overline{V_1})). \end{aligned}$$

So we may take $N = \overline{V_1}$. □

Remark 5.2. An examination of the proof of Lemma 5.1 shows that there is a variant of the lemma in which $\liminf$ is replaced by $\limsup$.

Proof of Theorem 1.1. (1) $\implies$ (2). Let ϕ be a pure state associated with $\text{Ind } \epsilon_z$ and $\mathcal{N}$ the usual weak*-neighbourhood base at zero in $(C_0(X) \times G)^*$. Suppose that $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) = r < k$. There exists $N \in \mathcal{N}$ such that

$$\liminf_n d((\text{Ind } \epsilon_{x_n}), \phi, N) = r$$

and hence a subsequence $(x_{n_i})_i$ such that

$$d((\text{Ind } \epsilon_{x_{n_i}}), \phi, N) = r < k \tag{5.1}$$

for all $i \geq 1$. Note that $(x_{n_i})_i$ converges k -times in X/G to z because $(x_n)_n$ does. The proof of [2, Theorem 2.3]¹ shows that, after passing to a further subsequence and reindexing,

¹The hypothesis in [2, Theorem 2.3] that all the orbits are locally closed in $\hat{A}$ is not needed in the case $A = C_0(X)$, since $\text{Ind } \epsilon_x$ is irreducible by [19, Proposition 4.2]. The representation $\tilde{\epsilon}_{x_{n_i}} \times \lambda$ on $L^2(G, \mu)$ used in [2, Theorem 2.3] is unitarily equivalent to our $\text{Ind } \epsilon_{x_{n_i}}$ on $L^2(G, \nu)$ via the unitary $W : L^2(G, \mu) \rightarrow L^2(G, \nu)$ defined by $(W\xi)(s) = \Delta(s)^{1/2}\xi(s)$.

there exist unit vectors $\eta_i^{(j)}$ ($1 \leq j \leq k$) in $L^2(G, \nu)$ such that

$$\lim_{i \rightarrow \infty} \langle \text{Ind } \epsilon_{x_{n_i}}(\cdot) \eta_i^{(j)}, \eta_i^{(j)} \rangle = \phi \quad (1 \leq j \leq k),$$

and that there exists i_0 such that if $i \geq i_0$ then $\eta_i^{(j)}$ and $\eta_i^{(l)}$ are orthogonal if $j \neq l$. By increasing i_0 if necessary, we may assume that $\langle \text{Ind } \epsilon_{x_{n_i}}(\cdot) \eta_i^{(j)}, \eta_i^{(j)} \rangle \in \phi + N$ for all $i \geq i_0$ and $1 \leq j \leq k$. Thus $d((\text{Ind } \epsilon_{x_{n_i}}), \phi, N) \geq k$ for all $i \geq i_0$, contradicting (5.1).

(2) $\implies$ (3). If $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \geq k$ then $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) > \lfloor k - \epsilon \rfloor$ for every $\epsilon > 0$. By Theorem 3.5, for every open neighbourhood V of z such that $\phi_z^{-1}(V)$ is relatively compact,

$$\nu(\phi_{x_n}^{-1}(V)) > (k - \epsilon)\nu(\phi_z^{-1}(V))$$

eventually, and hence (3) holds.

(3) $\implies$ (4) is immediate.

(4) $\implies$ (5). Let $(V_j)_{j \geq 1}$ be a decreasing sequence of basic open neighbourhoods of z such that $\phi_z^{-1}(V_1)$ is relatively compact (such neighbourhoods exist by Lemma 2.1). By Lemma 5.1 there exists a compact neighbourhood W_1 of z such that $W_1 \subset V_1$ and

$$\nu(\phi_{x_n}^{-1}(W_1)) > (k - 1)\nu(\phi_z^{-1}(W_1)).$$

Now assume there are compact neighbourhoods $W_1, W_2, \dots, W_m$ of z with $W_1 \supset W_2 \supset \dots \supset W_m$ such that

$$W_i \subset V_i \text{ and } \nu(\phi_{x_n}^{-1}(W_i)) > (k - 1)\nu(\phi_z^{-1}(W_i)) \quad (5.2)$$

for $1 \leq i \leq m$. Apply Lemma 5.1 to $(\text{Int } W_m) \cap V_{m+1}$ to obtain a compact neighbourhood W_{m+1} of z such that $W_{m+1} \subset (\text{Int } W_m) \cap V_{m+1}$ and (5.2) holds for $i = m + 1$.

(5) $\implies$ (1). We show first that $G \cdot x_n \rightarrow G \cdot z$ in X/G . Let $q : X \rightarrow X/G$ be the quotient map. Let U be a neighbourhood of $G \cdot z$ in X/G and $V = q^{-1}(U)$. There exists m such that $W_m \subset V$. Since $\liminf_n \nu(\phi_{x_n}^{-1}(W_m)) > 0$ there exists n_0 such that $\phi_{x_n}^{-1}(W_m) \neq \emptyset$ for $n \geq n_0$. Thus, for $n \geq n_0$,

$$G \cdot x_n = q(x_n) \in q(W_m) \subset q(V) = U.$$

Thus $G \cdot x_n$ is eventually in every neighbourhood of $G \cdot z$ in X/G .

Now suppose that $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) < \infty$. Then, as in the proof of Theorem 3.5, we may localise to an open G -invariant neighbourhood Y of z such that $G \cdot z$ is the unique limit in Y/G of the sequence $(G \cdot x_n)_n$. Eventually $W_m \subset Y$, and so the sequence $(x_n)_n$ converges k -times in Y/G to z by Proposition 4.1(1) applied to Y . But now $(x_n)_n$ converges k -times in X/G to z as well.

Finally, if $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) = \infty$ then $(x_n)_n$ converges k -times in X/G to z by Proposition 4.2. $\square$

Theorem 5.3. *Suppose that (G, X) is a free second-countable transformation group. Let $k \in \mathbb{P}$, let $z \in X$ and let $(x_n)_{n \geq 1} \subset X$ be a sequence such that $(G \cdot x_n)_n$ converges to $G \cdot z$ in X/G . Assume that $G \cdot z$ is locally closed in X . Then the following are equivalent:*

- (1) *there exists a subsequence $(x_{n_i})_{i \geq 1}$ of (x_n) which converges k -times in X/G to z ;*
- (2) $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \geq k$;

- (3) for every open neighbourhood V of z such that $\phi_z^{-1}(V)$ is relatively compact we have

$$\limsup_n \nu(\phi_{x_n}^{-1}(V)) \geq k\nu(\phi_z^{-1}(V));$$

- (4) there exists a real number $R > k - 1$ such that for every open neighbourhood V of z with $\phi_z^{-1}(V)$ relatively compact we have

$$\limsup_n \nu(\phi_{x_n}^{-1}(V)) \geq R\nu(\phi_z^{-1}(V));$$

- (5) there exists a decreasing sequence of basic compact neighbourhoods $(W_m)_{m \geq 1}$ of z such that, for each $m \geq 1$,

$$\limsup_n \nu(\phi_{x_n}^{-1}(W_m)) > (k - 1)\nu(\phi_z^{-1}(W_m)).$$

Proof. If (1) holds then $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_i}})) \geq k$ using Theorem 1.1, and hence

$$M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \geq M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_i}})) \geq M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_i}})) \geq k.$$

If (2) holds then by Lemma A.1 there is a subsequence $(x_{n_r})_r$ such that $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_r}})) = M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_r}}))$ so that $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_r}})) \geq k$. Let V be any open neighbourhood of z such that $\phi_z^{-1}(V)$ is relatively compact. Then

$$\limsup_n \nu(\phi_{x_n}^{-1}(V)) \geq \limsup_r \nu(\phi_{x_{n_r}}^{-1}(V)) \geq \liminf_r \nu(\phi_{x_{n_r}}^{-1}(V)) \geq k\nu(\phi_z^{-1}(V)),$$

where we have used Theorem 1.1 at the last step.

That (3) implies (4) is immediate. That (4) implies (5) follows from Remark 5.2 as in the proof of Theorem 1.1.

Assume (5). First suppose that $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) < \infty$. Since $G \cdot x_n \rightarrow_n G \cdot z$, we can localise to an open G -invariant neighbourhood Y of z such that $G \cdot z$ is the unique limit of $(G \cdot x_n)_n$, as in the proof of Theorem 3.5. Now (1) follows by applying Proposition 4.1(2) to Y .

If $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) = \infty$ then $(x_n)_n$ converges k -times in X/G to z by Proposition 4.2. $\square$

We now derive some further consequences of Theorem 1.1

Corollary 5.4. *Suppose that (G, X) is a free second-countable transformation group such that all the orbits are locally closed in X . Let $k \in \mathbb{P}$ and let $z \in X$. Then the following are equivalent:*

- (1) *there exists a sequence $(x_n)_{n \geq 1}$ in X which is k -times convergent in X/G to z ;*
- (2) *$M_U(\text{Ind } \epsilon_z) \geq k$.*

Proof. Assume (1). Then $M_U(\text{Ind } \epsilon_z) \geq M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \geq k$ by Theorem 1.1.

Assume (2). By [3, Lemma 1.2] there exists a sequence $(\pi_n)_{n \geq 1}$ converging to $\text{Ind } \epsilon_z$ such that $M_L(\text{Ind } \epsilon_z, (\pi_n)) = M_U(\text{Ind } \epsilon_z, (\pi_n)) = M_U(\text{Ind } \epsilon_z)$. Since the orbits are locally closed, the mapping $X \rightarrow (C_0(X) \times G)^\wedge : x \mapsto \text{Ind } \epsilon_x$ is a surjection. So there is a sequence $(x_n)_n$ in X such that $\text{Ind } \epsilon_{x_n} = \pi_n$ for each n . Then

$$M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) = M_L(\text{Ind } \epsilon_z, (\pi_n)) = M_U(\text{Ind } \epsilon_z) \geq k,$$

and it follows from Theorem 1.1 that $(x_n)_{n \geq 1}$ is k -times convergent in X/G to z . $\square$

That (1) $\implies$ (2) in Corollary 5.4 is a special case of [2, Theorem 2.3].

Corollary 5.5. *Suppose that (G, X) is a free second-countable transformation group such that all the orbits are locally closed in X . Let $k \in \mathbb{P}$ and let $z \in X$ such that $G \cdot z$ is not open in X . Then the following are equivalent:*

- (1) *whenever $(x_n)_{n \geq 1}$ is a sequence in X which converges to z and satisfies $z \notin G \cdot x_n$ eventually, then (x_n) is k -times convergent in X/G to z ;*
- (2) $M_L(\text{Ind } \epsilon_z) \geq k$.

Proof. Assume (1). By Lemma A.2, there is a sequence $(\pi_n)_{n \geq 1}$ in $(C_0(X) \times G)^\wedge$ such that $\pi_n \neq \text{Ind } \epsilon_z$ for all n , $\pi_n \rightarrow_n \text{Ind } \epsilon_z$ and

$$M_L(\text{Ind } \epsilon_z) = M_L(\text{Ind } \epsilon_z, (\pi_n)) = M_U(\text{Ind } \epsilon_z, (\pi_n)). \quad (5.3)$$

Since the orbits are locally closed, the mapping $X \rightarrow (C_0(X) \times G)^\wedge : x \mapsto \text{Ind } \epsilon_x$ is an open surjection. So there is a sequence $(x_n)_n$ in X such that $x_n \rightarrow z$ and $(\text{Ind } \epsilon_{x_n})_n$ is a subsequence of $(\pi_n)_n$. Using (5.3), $M_L(\text{Ind } \epsilon_z) = M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n}))$. By (1), $(x_n)_n$ is k -times convergent in X/G to z , so it follows from Theorem 1.1 that $M_L(\text{Ind } \epsilon_z) = M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) \geq k$.

Assume (2). If $(x_n)_n$ is a sequence in X which converges to z such that $z \notin G \cdot x_n$ eventually, then

$$M_L(\text{Ind } \epsilon_x, (\text{Ind } \epsilon_{x_n})) \geq M_L(\text{Ind } \epsilon_z) \geq k.$$

It follows from Theorem 1.1 that $(x_n)_n$ is k -times convergent in X/G to z . $\square$

Corollary 5.6 improves Proposition 4.2, and is immediate from Theorem 1.1.

Corollary 5.6. *Suppose that (G, X) is a free second-countable transformation group. Let $z \in X$ and let $(x_n)_{n \geq 1}$ be a sequence in X . Assume that $G \cdot z$ is locally closed in X . Then the following are equivalent:*

- (1) *for each $k \geq 1$, the sequence $(x_n)_n$ converges k -times in X/G to z ;*
- (2) $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) = \infty$;
- (3) *for every open neighbourhood V of z such that $\phi_z^{-1}(V)$ is relatively compact, $\nu(\phi_{x_n}^{-1}(V)) \rightarrow \infty$ as $n \rightarrow \infty$.*

Corollary 5.7. *Suppose that (G, X) is a free second-countable transformation group. Let $z \in X$ and let $(x_n)_{n \geq 1} \subset X$ be a sequence converging to z . Assume that $G \cdot z$ is locally closed in X . Then the following are equivalent:*

- (1) *there exists an open neighbourhood V of z such that $\phi_z^{-1}(V)$ is relatively compact and*

$$\limsup_n \nu(\phi_{x_n}^{-1}(V)) < \infty;$$

- (2) $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) < \infty$.

Proof. Suppose that (1) holds. Since $C_0(X) \times G$ is separable, it follows from Lemma A.1 that there exists a subsequence $(x_{n_j})_{j \geq 1}$ of (x_n) such that

$$M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_j}})) = M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_j}})) = M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})).$$

By (1) and Corollary 5.6, $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_j}})) < \infty$. Hence $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) < \infty$, that is (2) holds.

Suppose that (1) fails. Let $(V_i)_{i \geq 1}$ be a decreasing sequence of open basic neighbourhoods of z such that $\phi_z^{-1}(V_1)$ is relatively compact (such neighbourhoods exist by

Lemma 2.1). Then $\limsup_n \{\nu(\phi_{x_n}^{-1}(V_i))\} = \infty$ for each i and we may choose a subsequence $(x_{n_i})_i$ of $(x_n)_n$ such that $\nu(\phi_{x_{n_i}}^{-1}(V_i)) \rightarrow_i \infty$.

Let V be any open neighbourhood of z such that $\phi_z^{-1}(V)$ is relatively compact. There exists i_0 such that $V_i \subset V$ for all $i \geq i_0$. Then, for $i \geq i_0$,

$$\nu(\phi_{x_{n_i}}^{-1}(V_i)) \leq \nu(\phi_{x_{n_i}}^{-1}(V)).$$

Thus $\nu(\phi_{x_{n_i}}^{-1}(V)) \rightarrow_i \infty$. By Corollary 5.6, $M_L(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_{n_i}})) = \infty$. Hence $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_n})) = \infty$, that is (2) fails. $\square$

Corollary 5.8. *Suppose that (G, X) is a free second-countable transformation group such that all the orbits are locally closed in X . Let $z \in X$ and let $(x_n)_{n \geq 1} \subset X$ be a sequence converging to z . Then the following are equivalent:*

- (1) $M_U(\text{Ind } \epsilon_z) < \infty$;
- (2) *there exists an open neighbourhood V of z such that $\phi_z^{-1}(V)$ is relatively compact and*

$$\sup_{x \in V} \nu(\phi_x^{-1}(V)) < \infty;$$

Proof. If (2) holds then (1) holds by Corollary 3.7.

Let $(V_i)_{i \geq 1}$ be a decreasing sequence of open basic neighbourhoods of z such that $\phi_z^{-1}(V_1)$ is relatively compact. If (2) fails then $\sup_{x \in V_i} \{\nu(\phi_x^{-1}(V_i))\} = \infty$ for each i and we may choose a sequence $(x_i)_i$ such that $x_i \in V_i$ for all i and $\nu(\phi_{x_i}^{-1}(V_i)) \rightarrow_i \infty$. Since $(V_i)_i$ is decreasing, $x_i \rightarrow_i z$.

Let V be an open neighbourhood of z such that $\phi_z^{-1}(V)$ is relatively compact. There exists i_0 such that $V_i \subset V$ for all $i \geq i_0$. Then, for $i \geq i_0$,

$$\nu(\phi_{x_i}^{-1}(V_i)) \leq \nu(\phi_{x_i}^{-1}(V)).$$

Thus $\nu(\phi_{x_i}^{-1}(V)) \rightarrow_i \infty$. By the contrapositive of the (2) $\implies$ (1) direction of Corollary 5.7, $M_U(\text{Ind } \epsilon_z, (\text{Ind } \epsilon_{x_i})) = \infty$. Hence $M_U(\text{Ind } \epsilon_z) = \infty$, that is (1) fails. $\square$

In the terminology of [13], property (2) of Corollary 5.8 says that V is an integrable neighbourhood of z ; in view of [5, Theorem 2.8], the equivalence of (1) and (2) is essentially contained in [13, Theorem 5.8(2)].

Remark 5.9. One can formulate conditions equivalent to 4.2 (2), 5.6 (3), 5.8 (2) and 5.7 (1), respectively, by suppressing the requirement that $\phi_z^{-1}(V)$ is relatively compact. This is because $G \cdot z$ is assumed to be locally closed in X and hence every neighbourhood V of z contains a neighbourhood V_1 of z such that $\phi_z^{-1}(V_1)$ is relatively compact (see Lemma 2.1).

6. EXAMPLES

We apply our results in two examples. First, we calculate upper and lower multiplicities relative to subsequences in the transformation group described by Rieffel in [18, Example 1.18]. Then we combine the ideas of [18, Example 1.18] and [16, Example on p. 298] to construct a transformation group with a sequence $(G \cdot x_n)_n$ of orbits converging to two

distinct orbits $G \cdot x_0$ and $G \cdot z_0$ such that

$$\begin{aligned} M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) &= M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = 2 \\ M_U(\text{Ind } \epsilon_{z_0}, (\text{Ind } \epsilon_{x_n})) &= M_L(\text{Ind } \epsilon_{z_0}, (\text{Ind } \epsilon_{x_n})) = 3. \end{aligned}$$

Example 6.1. In [18, Example 1.18], the space X is a closed subset of $\mathbb{R}^3$ and the group is $G = \mathbb{R}$. The action is free, with all orbits closed in X . The orbit space X/G is a compact Hausdorff space, and is discrete except for one limit point. This limit point is $\{(0, s, 0) : s \in \mathbb{R}\}$ with the action of $\mathbb{R}$ on it being translation, and orbit representative $x_0 = (0, 0, 0)$.

Let $(b_n)_{n \geq 1}$ be a strictly decreasing sequence of real numbers converging to 0. If $n \geq 1$ then the n th orbit representative is $x_n = (b_n, 0, 0)$, and assigned to each orbit is a repetition integer $L_n \geq 0$.

If $L_n = 0$ then the orbit of x_n is the line $\{(b_n, s, 0) : s \in \mathbb{R}\}$ with the action of $\mathbb{R}$ on it being translation. If $L_n \geq 1$, choose a strictly decreasing finite sequence $(b_n^j)_j$ of length $L_n + 1$, with $b_{n+1} < b_n^{L_n} < \dots < b_n^2 < b_n^1 < b_n^0 = b_n$. Let $x_n^j = (b_n^j, 0, 0)$ for $0 \leq j \leq L_n$; note that $x_n = x_n^0$. The points x_n^j ($j = 0, \dots, L_n$) are in the orbit $G \cdot x_n$. The n th orbit consists of $L_n + 1$ vertical line segments parallel to the y -axis, each of these line segments goes through x_n^j for some j . The vertical line segments are joined by L_n arcs. The action is described by specifying what happens to the orbit representatives x_n ($n \geq 1$):

$$t \cdot x_n = \begin{cases} (b_n, t, 0) & \text{if } t \in (-\infty, n]; \\ (b_n^{L_n}, s, 0) & \text{if } s \in (-n, \infty) \text{ and } t = s + L_n(2n + 1); \\ (b_n^j, s, 0) & \text{if } s \in (-n, n] \text{ and } t = s + j(2n + 1) \\ & \text{for } j \in \{1, \dots, L_n - 1\}; \\ ((1 - s)b_n^j + sb_n^{j+1}, n \cos(\pi s), n \sin(\pi s)) & \text{if } s \in (0, 1] \text{ and } t = s + n + j(2n + 1), \\ & \text{for } j \in \{0, \dots, L_n - 1\}. \end{cases}$$

Lemma 6.2. *Let (G, X) be the free transformation group from Example 6.1 with orbit representatives x_n . Then*

$$\begin{aligned} M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) &= \liminf_n (L_n + 1); \\ M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) &= \limsup_n (L_n + 1). \end{aligned}$$

Proof. Suppose that $\liminf_n (L_n + 1) = k < \infty$. We will show that $M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) \geq k$ and $M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) \leq k$ using Theorems 1.1 and 3.5.

Since $\liminf_n (L_n + 1) = k$ there exists n_0 such that $L_n + 1 \geq k$ for $n \geq n_0$. For each $n \geq n_0$ and $j \in \{0, \dots, k - 1\}$ choose $t_n^{(j)} = j(2n + 1)$ for . Then, as $n \rightarrow \infty$, we have

$$\begin{aligned} t_n^{(0)} \cdot x_n &= x_n \rightarrow x_0 \\ t_n^{(j)} \cdot x_n &= (b_n^j, 0, 0) = q_n^j \rightarrow x_0 \quad \text{for } j \in \{1, \dots, k - 1\} \\ t_n^{(i)} - t_n^{(j)} &= (i - j)(2n + 1) \rightarrow \infty \quad \text{if } i \neq j. \end{aligned}$$

So $(x_n)_n$ converges k -times in X/G to x_0 , and hence $M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) \geq k$ by Theorem 1.1.

Now consider the (relatively) open relatively compact neighbourhood

$$V = ([0, 1) \times (-1/2, 1/2) \times (-1/2, 1/2)) \cap X \tag{6.1}$$

of x_0 . Since $\liminf_n (L_n + 1) = k$ there exists a subsequence $(x_{n_i})_i$ such that $L_{n_i} + 1 = k$ and $x_{n_i} \in V$ for all i . Note that the arcs of $G \cdot x_{n_i}$ joining the parallel line segments of the orbit do not meet V . So if $t \cdot x_{n_i} \in V$ then t is either in the interval $(-1/2, 1/2)$ or in a translate by $j(2n_i + 1)$ of it, where $1 \leq j \leq k - 1$. Thus $\phi_{x_0}^{-1}(V) = (-1/2, 1/2)$ is relatively compact and

$$\nu(\phi_{x_{n_i}}^{-1}(V)) = k\nu(\phi_{x_0}^{-1}(V)) < \infty$$

for all i , so $M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) \leq k$ by Theorem 3.5. Thus $M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = k$

If $\liminf_n (L_n + 1) = \infty$, then for any $k \in \mathbb{P}$ we have $L_n + 1 \geq k$ eventually, and $(x_n)_n$ converges k -times to x_0 in X/G as above. Thus $M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = \infty$.

Now suppose that $\limsup_n (L_n + 1) = k < \infty$. Then there exists a subsequence $(x_{n_i})_i$ such that $L_{n_i} + 1 = k$ for all i . As above, (x_{n_i}) converges k -times to x_0 in X/G , and hence $M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) \geq k$ by Theorem 5.3. On the other hand, there exists n_1 such that for all $n \geq n_1$, $x_n \in V$ and $L_n + 1 \leq k$. We have

$$\nu(\phi_{x_n}^{-1}(V)) = k\nu(\phi_{x_0}^{-1}(V)) = k < \infty$$

whenever $n \geq n_1$, and hence $M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) \leq k$ by Theorem 3.6. Thus $M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = k$.

If $\limsup_n (L_n + 1) = \infty$ then, given $k \in \mathbb{P}$, there exists a subsequence $(x_{n_i})_i$ such that $L_{n_i} + 1 \geq k$. Then (x_{n_i}) converges k -times to x_0 in X/G and $M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) \geq k$. Thus $M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = \infty$. $\square$

Remark 6.3. In the situation of Lemma 6.2 one can easily show that $M_L(\text{Ind } \epsilon_{x_0}) = M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n}))$ and $M_U(\text{Ind } \epsilon_{x_0}) = M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n}))$ using Lemma A.2 and [3, Lemma 1.2], respectively.

Example 6.4. The idea of this example is to splice together two instances of Example 6.1, one with constant repetition integer 1 and the other with constant repetition integer 2.

Again, the space X is a closed subspace of $\mathbb{R}^3$, the action of the group $G = \mathbb{R}$ is free and the orbits are closed in X . The orbit space is compact but non-Hausdorff. This time we have two limit orbits, $\{(0, s, 0) : s \in \mathbb{R}\}$ and $\{(1, s, 0) : s \in \mathbb{R}\}$ with orbit representatives $x_0 = (0, 0, 0)$ and $z_0 = (1, 0, 0)$; the action of $\mathbb{R}$ on these two orbits is by translation.

If $n \geq 1$, the data for the n th orbit is:

$$b_n = 2^{-2n}, \quad b_n^1 = 2^{-(2n+1)}, \quad a_n = 1 - 2^{-3n}, \quad a_n^1 = 1 - 2^{-(3n+1)}, \quad a_n^2 = 1 - 2^{-(3n+2)};$$

the orbit representative is

$$x_n = \left(\frac{a_n + b_n}{2}, 0, n \right).$$

The orbit consists of five vertical line segments parallel to the y -axis, two on the left of and three on the right of the line $\{(1/2, s, 0) : s \in \mathbb{R}\}$; the line segments are joined by four arcs. Again we describe the action by specifying what happens to the orbit representatives

x_n . If $n \geq 1$ and $u \in \mathbb{R}$ then

$$u \cdot x_n \left\{ \begin{array}{ll} (b_n^1, u - 3n - \frac{3}{2}, 0) & \text{if } u \in [2n + \frac{3}{2}, \infty); \\ (b_n + s(b_n^1 - b_n), n \cos(\pi s), n \sin(\pi s)) & \text{if } u \in [2n + \frac{1}{2}, 2n + \frac{3}{2}) \\ & \text{and } s = u - 2n - \frac{1}{2}; \\ (b_n, u - n - \frac{1}{2}, 0) & \text{if } u \in [\frac{1}{2}, 2n + \frac{1}{2}); \\ (a_n + (u + \frac{1}{2})(b_n - a_n), -n \sin(u\pi), n \cos(u\pi)) & \text{if } u \in [-\frac{1}{2}, \frac{1}{2}); \\ (a_n, u + n + \frac{1}{2}, 0) & \text{if } u \in [-2n - \frac{1}{2}, -\frac{1}{2}); \\ (a_n^1 + s(a_n - a_n^1), n \cos(\pi s), n \sin(\pi s)) & \text{if } u \in [-2n - \frac{3}{2}, -2n - \frac{1}{2}) \\ & \text{and } s = u + 2n + \frac{3}{2}; \\ (a_n^1, u + 3n + \frac{3}{2}, 0) & \text{if } u \in [-4n - \frac{3}{2}, -2n - \frac{3}{2}); \\ (a_n^2 + s(a_n^1 - a_n^2), n \cos(\pi s), n \sin(\pi s)) & \text{if } u \in [-4n - \frac{5}{2}, -4n - \frac{3}{2}), \\ & \text{and } s = u + 4n + \frac{5}{2}; \\ (a_n^2, u + 5n + \frac{5}{2}, 0) & \text{if } u \in (-\infty, -4n - \frac{5}{2}). \end{array} \right.$$

It is straightforward to verify that $M_U(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = M_L(\text{Ind } \epsilon_{x_0}, (\text{Ind } \epsilon_{x_n})) = 2$ and $M_U(\text{Ind } \epsilon_{z_0}, (\text{Ind } \epsilon_{x_n})) = M_L(\text{Ind } \epsilon_{z_0}, (\text{Ind } \epsilon_{x_n})) = 3$ using the methods of Lemma 6.2

APPENDIX A.

The purpose of this appendix is to establish sequence versions of [6, Propositions 2.2 and 2.3] which were used in §3 and §5. See §2 for the relevant notation.

Lemma A.1. *Let A be a separable C^* -algebra, $\pi \in \hat{A}$ and $(\pi_n)_{n \geq 1}$ a sequence in $\hat{A}$. Then there exists a subsequence $(\pi_{n_i})_i$ such that*

$$M_L(\pi, (\pi_{n_i})_i) = M_U(\pi, (\pi_{n_i})_i) = M_U(\pi, (\pi_n)).$$

Proof. Let ϕ be a pure state of A associated with π . Suppose that $M_U(\pi, (\pi_n)) = m$ (where possibly $m = \infty$). It suffices to construct (π_{n_i}) such that $M_L(\pi, (\pi_{n_i})_i) \geq m$. By assumption, there exists $N_0 \in \mathcal{N}$ such that, for $N \in \mathcal{N}$ satisfying $N \subset N_0$,

$$M_U(\phi, N, (\pi_n)) = \limsup_n d(\pi_n, \phi, N) = m. \quad (\text{A.1})$$

Since A is separable, the weak*-topology on A^* is first countable and so there is a basic decreasing sequence $(N_i)_{i \geq 1}$ in $\mathcal{N}$ such that $N_i \subset N_0$ for all $i \geq 1$. By (A.1), for all $i \geq 1$,

$$\limsup_n d(\pi_n, \phi, N_i) = m. \quad (\text{A.2})$$

Firstly, suppose that $m < \infty$. By (A.2), we may construct a strictly increasing sequence $(n_i)_{i \geq 1}$ such that $d(\pi_{n_i}, \phi, N_i) = m$ for all $i \geq 1$. Let $N \in \mathcal{N}$. There exists i_0 such that $N_{i_0} \subset N$. For $i \geq i_0$,

$$d(\pi_{n_i}, \phi, N) \geq d(\pi_{n_i}, \phi, N_i) = m.$$

Thus

$$M_L(\phi, N, (\pi_{n_i})) = \liminf_i d(\pi_{n_i}, \phi, N) \geq m$$

and hence

$$M_L(\pi, (\pi_{n_i})) = \inf_{N \in \mathcal{N}} M_L(\phi, N, (\pi_{n_i})) \geq m.$$

Secondly, suppose that $m = \infty$. By (A.2), we may construct a strictly increasing sequence $(n_i)_{i \geq 1}$ such that $d(\pi_{n_i}, \phi, N_i) \geq i$ for all $i \geq 1$. Let $N \in \mathcal{N}$. There exists i_0 such that $N_{i_0} \subset N$. For $i \geq i_0$,

$$d(\pi_{n_i}, \phi, N) \geq d(\pi_{n_i}, \phi, N_i) \geq i.$$

Thus

$$M_L(\phi, N, (\pi_{n_i})) = \liminf_i d(\pi_{n_i}, \phi, N) = \infty$$

and hence

$$M_L(\pi, (\pi_{n_i})) = \inf_{N \in \mathcal{N}} M_L(\phi, N, (\pi_{n_i})) = \infty.$$

□

Lemma A.2. *Let A be a separable C^* -algebra. Let $\pi \in \hat{A}$ such that $\{\pi\}$ is not open in $\hat{A}$. Then there exists a sequence $(\pi_n)_{n \geq 1}$ in $\hat{A}$ such that $\pi_n \neq \pi$ for all $n \geq 1$, $\pi_n \rightarrow_n \pi$, and*

$$M_L(\pi) = M_L(\pi, (\pi_n)) = M_U(\pi, (\pi_n)).$$

Proof. First assume that $M_L(\pi) = \infty$. Since $\hat{A}$ is second countable and $\{\pi\}$ is not open in $\hat{A}$, there exists a sequence $(\pi_n)_{n \geq 1}$ in $\hat{A}$ such that $\pi_n \rightarrow \pi$ and $\pi_n \neq \pi$ for all n . Then

$$\infty = M_L(\pi) \leq M_L(\pi, (\pi_n)) \leq M_U(\pi, (\pi_n)),$$

and we must have equality throughout.

Next, assume that $M_L(\pi) = k$ where $1 \leq k < \infty$. Let ϕ be a pure state associated with π . There exists $N_0 \in \mathcal{N}$ such that if $N \in \mathcal{N}$ and $N \subset N_0$, then $M_L(\phi, N) = \liminf_{\sigma \rightarrow \pi, \sigma \neq \pi} d(\sigma, \phi, N) = k$. Since A is separable, there exists a decreasing base $(V_n)_{n \geq 1}$ of open neighbourhoods of π in $\hat{A}$. We have

$$\liminf_{\sigma \rightarrow \pi, \sigma \neq \pi} d(\sigma, \phi, N_0) = k,$$

so for $n \geq 1$ there exist $\pi_n \in V_n \setminus \{\pi\}$ such that $d(\pi_n, \phi, N_0) = k$. Since $\{V_n\}$ is decreasing, $\pi_n \rightarrow \pi$ as $n \rightarrow \infty$. Note that

$$\begin{aligned} M_U(\pi, (\pi_n)) &= \inf_{N \in \mathcal{N}} M_U(\phi, N, (\pi_n)) \\ &\leq M_U(\phi, N_0, (\pi_n)) \\ &= \limsup_n d(\pi_n, \phi, N_0) = k. \end{aligned}$$

Now $k = M_L(\pi) \leq M_L(\pi, (\pi_n)) \leq M_U(\pi, (\pi_n)) \leq k$ and the result follows. □

REFERENCES

- [1] R.J. Archbold, *Upper and lower multiplicity for irreducible representations of C^* -algebras*, Proc. London Math. Soc. **69** (1994), 121–143.
- [2] R.J. Archbold and K. Deicke, *Bounded trace C^* -algebras and integrable actions*, Math. Zeit. **250** (2005), 393–410.
- [3] R.J. Archbold and E. Kaniuth, *Upper and lower multiplicity for irreducible representations of SIN-groups*, Illinois J. Math. **43** (1999), 692–706.
- [4] R.J. Archbold, E. Kaniuth, J. Ludwig, G. Schlichting and D.W.B. Somerset, *Strength of convergence in duals of C^* -algebras and nilpotent Lie groups*, Adv. Math. **158** (2001), 26–65.
- [5] R.J. Archbold, D.W.B. Somerset and J.S. Spielberg, *Upper multiplicity and bounded trace ideals in C^* -algebras*, J. Funct. Anal. **146** (1997), 430–463.

- [6] R.J. Archbold and J.S. Spielberg, *Upper and lower multiplicity for irreducible representations of C^* -algebras. II*, J. Operator Theory **36** (1996), 201–231.
- [7] K. Deicke, *Pointwise unitary coactions on C^* -algebras with continuous trace*, Dissertation, Universität Paderborn, 1999.
- [8] J. Dixmier, *C^* -algebras*, North-Holland, Paris 1969.
- [9] J. Glimm, *Locally compact transformation groups*, Trans. Amer. Math. Soc. **101** (1961), 124–138.
- [10] E.C. Gootman, *The type of some C^* - and W^* -algebras associated with transformation groups*, Pacific J. Math. **48** (1973), 93–106.
- [11] P. Green, *C^* -algebras of transformation groups with smooth orbit space*, Pacific J. Math. **72** (1977), 71–97.
- [12] A. an Huef, *The transformation groups whose C^* -algebras are Fell algebras*, Bull. London Math. Soc. (2000), 73–76.
- [13] A. an Huef, *Integrable actions and the transformation groups whose C^* -algebras have bounded trace*, Indiana Univ. Math. J. **51** (2002), 1197–1233.
- [14] J.L. Kelley, *General Topology*, Van Nostrand Reinhold, New York, 1970.
- [15] J. Ludwig, *On the behaviour of sequences in the dual of a nilpotent Lie group*, Math. Ann. **287** (1990), 239–257.
- [16] R.S. Palais, *On the existence of slices for actions of non-compact Lie groups*, Ann. of Math. **73** (1961), 295–323.
- [17] G. K. Pedersen, *Analysis Now*, Springer, New York, 1989.
- [18] M.A. Rieffel, *Integrable and proper actions on C^* -algebras, and square-integrable representations of groups*, Expo. Math. **22** (2004), 1–53.
- [19] D.P. Williams, *The topology on the primitive ideal space of transformation group C^* -algebras and CCR transformation group C^* -algebras*, Trans. Amer. Math. Soc. **266** (1981), 335–359.
- [20] D.P. Williams, *Transformation group C^* -algebras with continuous trace*, J. Funct. Anal. **41** (1981), 40–76.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ABERDEEN, ABERDEEN AB24 3UE,
SCOTLAND, UNITED KINGDOM

E-mail address: r.archbold@maths.abdn.ac.uk

SCHOOL OF MATHEMATICS, UNIVERSITY OF NEW SOUTH WALES, SYDNEY, NSW 2052, AUSTRALIA

E-mail address: astrid@unsw.edu.au